arXiv:1611.01741v1 [math.GR] 6 Nov 2016

ON COMMENSURABILITY OF SOME RIGHT-ANGLED ARTIN GROUPS

MONTSERRAT CASALS-RUIZ, ILYA KAZACHKOV, AND ALEXANDER ZAKHAROV

Abstract. In this paper we study the classification of right-angled Artin groups up to commensurability. We characterise the commensurability classes of RAAGs defined by trees of diameter 4. In particular, we prove a conjecture of Behrstock and Neumann that there are infinitely many commensurability classes. Hence, we give first examples of RAAGs that are quasi-isometric but not commensurable.

1. Introduction

1.1. **Context.** One of the basic problems on locally compact topological groups is to classify their lattices up to commensurability. Recall that two lattices $\Gamma_1, \Gamma_2 < G$ are commensurable if and only if there exists $g \in G$ such that $\Gamma_1 \cap \Gamma_2^g$ has finite index in both Γ_1 and Γ_2^g. In particular, commensurable lattices have covolumes that are commensurable rational numbers, that is, they have a rational ratio.

The notion of commensurability was generalized to better suit topological and large-scale geometric properties and to compare groups without requiring them to be subgroups of a common group. More precisely, we say that two groups H and K are (abstractly) commensurable if they have isomorphic finite index subgroups. In this article, we will only be concerned with the notion of abstract commensurability and we simply refer to it as commensurability.

As we mentioned, commensurability is closely related to the large-scale geometry of the group. Indeed, any finitely generated group can be endowed with a natural word-metric which is well-defined up to quasi-isometry and since any finitely generated group is quasi-isometric to any its finite index subgroup, it follows that commensurable groups are quasi-isometric.

Gromov suggested to study groups from this geometric point of view and understand the relation between these two concepts. More precisely, a basic problem in geometric group theory is to classify commensurability and quasi-isometry classes (perhaps within a certain class) of finitely generated groups and to understand whether or not these classes coincide.

The classification of groups up to commensurability (both in the abstract and classical case) has a long history and a number of famous solutions for very diverse classes of groups such as Lie groups and more generally, locally compact topological groups, hyperbolic 3-manifold groups, pro-finite

Key words and phrases. right-angled Artin groups, commensurability, quasi-isometries.

This work was supported by the ERC Grant 336983, by the Basque Government grant IT974-16 and by the grant MTM2014-53810-C2-2-P of the Ministerio de Economia y Competitividad of Spain. The first author was supported by Juan de la Cierva programme of the Spanish Government. The third author was supported by the Russian Foundation for Basic Research (project no. 15-01-05823).

groups, Grigorchuk-Gupta-Sidki groups, etc see for instance [BJN09, DW93, Mar73, D85, Si43, GrW03, Ga16].

In this paper, we focus on the question of classification of right-angled Artin groups, RAAGs for short, up to commensurability. Recall that a RAAG is a finitely presented group $G(\Gamma)$ which can be described by a finite simplicial graph Γ, the commutation graph, in the following way: the vertices of Γ are in bijective correspondence with the generators of $G(\Gamma)$ and the set of defining relations of $G(\Gamma)$ consists of commutation relations, one for each pair of generators connected by an edge in Γ.

RAAGs have become central in group theory, their study interweaves geometric group theory with other areas of mathematics. This class interpolates between two of the most classical families of groups, free and free abelian groups, and its study provides uniform approaches and proofs, as well as rich generalisations of the results for free and free abelian groups. The study of this class from the different perspectives has contributed to the development of new, rich theories such as the theory of CAT(0) cube complexes and has been an essential ingredient in Agol's solution to the virtually fibred Conjecture.

The commensurability classification of RAAGs has been previously considered for the following classes of RAAGs:

- Free groups [St68, D85, KPS73], [Gr93, 1.C];
- Free Abelian groups, [Gr81, B72];
- $F_m \times Z^n$, [Wh10];
- Free products of free groups and free Abelian groups, [BJN09];
- $F_m \times F_n$ with $m, n \geq 2$, [Wi96, BM00];
- $G(\Gamma)$, where Γ is a tree of diameter ≤ 3;
- $G(\Gamma)$, when the outer automorphism of G is finite, Γ is star-rigid and does not have induced 4-cycles, [H16].

It turns out that, inside the class of RAAGs, the classification up to commensurability coincides with the quasi-isometric classification for all known cases. These rigidity results are mainly a consequence of the rigid structure of the intersection pattern of flats in the universal cover of the Salvetti complex.

In this paper we describe the commensurability classes of RAAGs defined by trees of diameter at most 4 and describe the "minimal" group in each commensurability class (minimal in terms of number of generators or the rank of its abelianization). In particular, we show that there exist infinitely many different commensurability classes confirming a conjecture of Behrstock and Neumann. In their paper [BN08], the authors show that RAAGs defined by trees of diameter at least 3 are quasi-isometric, so we provide first examples of RAAGs that are quasi-isometric but not commensurable. As in the classical case of lattices in locally compact topological groups, we define an ordered set that plays the role of the covolume and prove that the groups are commensurable if and only if the sets are commensurable, that is they have the same cardinality and constant rational ordered ratios.

1.2. **Main results.** Let $\Delta = (V(\Delta), E(\Delta))$ be a graph without loops and double edges, then we denote by $\mathbb{G}(\Delta)$ the RAAG defined by the commutation graph Δ. We call the vertices of the graph Δ the *canonical generators* of $\mathbb{G}(\Delta)$.

For our purposes, it will be convenient to encode finite trees of diameter four as follows. Let T be any finite tree of diameter four. Let f be a path (without backtracking) of length four from one leaf of T to another. By definition f contains 5 vertices and let $c_f \in V(T)$ be the middle vertex in f. It is immediate to see that the choice of the vertex $c = c_f$ does not depend on the choice of the path f of length four. We call c the *center* of T.

Any leaf of T connected to c by an edge is called a *hair vertex*. Vertices connected to c by an edge which are not hair are called *pivots*. Any finite tree T of diameter 4 is uniquely defined by the number q of hair vertices and by the number k_i of pivots of a given degree $d_i + 1$. Hence we encode any tree of diameter 4 as $T((d_1, k_1), \dots, (d_l, k_l); q)$. Here all d_i and k_i and l are positive integers, q is a non-negative integer, and all d_i are different; moreover, either $l \geq 2$ or $l = 1$ and $k_1 \geq 2$, so that T indeed has diameter 4. See Figure 1.



FIGURE 1. Tree of diameter 4

Given a tree of diameter four $T = T((d_1, k_1), \dots, (d_l, k_l); q)$, we denote by $M(T) = M(T((d_1, k_1), \dots, (d_l, k_l); q))$ the set of different numbers d_i, that is, renaming if necessary, we have that $M(T((d_1, k_1), \dots, (d_l, k_l); q))) = \{d_1 < d_2 < \cdots < d_l\}$.

In the original usage two rational numbers $a, b \in \mathbb{R}$ are commensurable if and only if the ratio $\frac{a}{b}$ is rational. In this fashion, we will say that two ordered sets $P = \{p_1 < \cdots < p_k\}$ and $Q = \{q_1 < \cdots < q_l\}$ are *commensurable* if they have the same cardinality, i.e. $k = l$, and each quotient $\frac{p_i}{q_i} = c \in \mathbb{Q}$. In this case we write $P = cQ$.

We say that a set $M = \{m_1, \dots, m_k\}$, $m_i \in \mathbb{N}$, is minimal if the greatest common divisor $d = \gcd(m_1, \dots, m_k)$ is 1. It is clear that in each commensurability class of a set $M \subset \mathbb{N}^k$, there exists a minimal set that belongs to the class, namely $\{\frac{m_1}{d}, \dots, \frac{m_k}{d}\}$.

We show that the commensurability class of the set $M(T((d_1,k_1),\dots,(d_l,k_l);q))$ determines the commensurability class of the RAAG defined by the tree of a diameter 4.

Theorem 6 (Characterisation of commensurability classes)**.** *Let T and T' be two finite trees of diameter 4, $T=T((d_1,k_1),\dots,(d_l,k_l);q)$ and $T'=T((d'_1,k'_1),\dots,(d'_{l'},k'_{l'});q')$. Let $G=\mathbb{G}(T)$ and $G'=\mathbb{G}(T')$. Consider the sets $M=M(T)$, $M'=M(T')$. Then G and G' are commensurable if and only if M and M' are commensurable.*

For $n>1$ denote by P_n the path with n vertices and $n-1$ edges. By a minimal RAAG in some class C we mean a RAAG in C with the minimal number of generators, i.e., defined by a graph with minimal number of vertices among all commutation graphs of RAAGs in C.

Theorem 7. *Let $T=T((d_1,k_1),\dots,(d_l,k_l);q)$ be a finite tree of diameter 4. Let $\mathcal{C}(T)$ be the commensurability class of $\mathbb{G}(T)$ and let $M=M(T)$ be as above, so $|M|=l$. Then the minimal RAAG that belongs to $\mathcal{C}(T)$ is either the RAAG defined by the tree $T'=T((d'_1,1),\dots,(d'_l,1);0)$, where $M(T')$ is minimal in the commensurability class of M, if $|M|>1$, or the RAAG defined by the path of diameter 3, that is $\mathbb{G}(P_4)$, if $|M|=1$.*

Our results extend naturally to the commensurability classification of some right-angled Coxeter groups. Recall that every RAAG G embeds naturally as a finite index subgroup into a right-angled Coxeter group, say $C(G)$. Hence, we have the following result

Theorem 1. *There are infinitely many pair-wise quasi-isometric, but pair-wise not commensurable right-angled Coxeter groups defined by graphs of diameter 4 with cliques of dimension* 2.

In their work [BJN10], Behrstock and Neumann introduced the notion of n-trees and gave a description of the quasi-isometry classes of RAAGs defined by n-trees. In particular, to each n-tree, the authors associate a 2-coloured graph, which corresponds to a graph of groups decomposition, where vertex groups are isomorphic to $Z^{n-1}\times F_v$ and edge groups are isomorphic to Z^n. As trees of diameter 4 have a central vertex, n-trees have a central $n-1$-simplex, say $(c_1,\dots,c_{n-1})$. Vertices of degree $n-1$ are analogs of leaves of a tree. If a "leaf" w on the n-tree only belongs to the n-simplex $(c_1,\dots,c_{n-1},w)$, we say that w is a hair vertex. Otherwise, vertices which do not belong to the central simplex and are not "leaves" of the n-tree are called pivots of the n-tree. Hence, one can encode an n-tree nT of diameter 4 (i.e. those n-trees whose associated graph has diameter 4) as a tuple $((n_1,m_1),\dots,(n_l,m_l);s)$, where m_i is the rank of the pivot minus 1. As above, let $M(nT)=\{m_1,\dots,m_l\}$.

The interested reader can check that the proofs given in this paper for paths of diameter 4 extend almost verbatim to n-trees of diameter 4 and we have the following

Theorem 2. *Let nT and nT' be two n-trees of diameter* 4*. Then $\mathbb{G}(nT)$ and $\mathbb{G}(nT')$ are commensurable if and only if the set $M(nT)$ and $M(nT')$ are commensurable.*

1.3. **Strategy of the proof.** As we discussed, in previous results on commensurability of RAAGs, the structure of the intersection pattern of flats inside the universal cover of the Salvetti complex is so rigid that the large-scale geometry that it determines forces commensurability.

In our case, all trees of diameter 4 are quasi-isometric and so geometry is not sufficient to determine the commensurability classes. However, we will use the structure of the intersection pattern of flats together with algebra to derive the result.

We encode the intersection pattern in a finite graph obtained as a quotient of the reduced extension graph of the RAAG by the action of the subgroup H. This finite graph has an abstract algebraic interpretation: its vertices correspond to conjugacy classes of cyclic subgroups whose centraliser is non-abelian; and its edges are defined via commutation of some representatives in the class. Defined this way, the graph is an algebraic invariant, that is if $\varphi : H \to H'$ is an isomorphism, then φ induces an isomorphism between the finite graphs associated to H and H' and so an isomorphism between the (conjugacy classes of) non-abelian centralisers. In our case, non-abelian centralisers are of the form $\mathbb{Z} \times F_n$ and so we can finally obtain isomorphism between free groups.

The isomorphism between subgroups of free groups of different ranks imposes conditions on the indices of the corresponding subgroups, and so, restrictions on the powers of conjugates of generators that belong to H and H'. We encode these conditions in a linear system of equations. If the subgroups H and H' are isomorphic, then the system must have positive integer solutions. Finally, we show that for the system to have positive solutions the sets $M(T)$ and $M(T')$ need to be commensurable.

We believe that the general strategy of our proof, that is to reduce the existence of subgroups to a linear system of equations with positive integer solutions, can be used to study commensurability classes of RAAGs defined by trees and more general RAAGs. The real obstacle is to elucidate the necessary conditions for the system of equations to have positive integer solutions – "just" a linear algebra problem.

2. Necessary conditions for commensurability

Below we always conjugate as follows $g^h = hgh^{-1}$.

Note that centralisers in RAAGs are isolated, that is for any element $g \in \mathbb{G}(\Delta)$ and any non-zero integer m we have that $C_G(g) = C_G(g^m)$, see [Ser89].

Recall that for any RAAG given by the commutation graph Δ the *extension graph* Δ^e is defined as follows, see [KK13]. The vertex set of Δ^e is the set of all elements of $\mathbb{G}(\Delta)$ which are conjugate to the canonical generators (vertices of Δ). Two vertices are joined by an edge if and only if the corresponding group elements commute. The group $\mathbb{G}(\Delta)$ acts on Δ^e by conjugation.

In some sense, the extension graph encodes the structure of the intersection pattern of flats inside the universal cover of the Salvetti complex. It plays an essential role in establishing the quasi-isometric rigidity of the class of RAAGs with finite outer automorphism groups, see [H16].

Observe that if the graph Δ is a tree, then $\mathbb{G}(\Delta)$ splits as the fundamental group of a graph of groups as follows. The underlying graph is Δ, and the vertex group at every vertex is defined to be the centralizer of the corresponding canonical generator. Note that if v is some vertex of Δ, and $u_1, \dots, u_s$ are all vertices of Δ adjacent to v, then $C(v) = \langle v, u_1, \dots, u_s \rangle \cong \mathbb{Z} \times F_s$, where F_s is the free group of rank s. In particular, $C(v)$ is abelian if and only if v has degree 1, and in this case $C(v) \cong \mathbb{Z}^2$ is contained in the centralizer of the vertex adjacent to v. For an edge e connecting vertices u and v the edge group at e is $C(u) \cap C(v) = \langle u, v \rangle \cong \mathbb{Z}^2$. We call this splitting the *centralizer splitting* of $\mathbb{G}(\Delta)$.

Note that the centralizer splitting is neither reduced nor minimal, since for every vertex of degree 1 in Δ the vertex group is equal to the incident edge group. Thus it makes sense to consider the

reduced centralizer splitting of $\mathbb{G}(\Delta)$ (for a tree Δ), which is obtained from the centralizer splitting by removing all vertices of degree 1. It is easy to see that this splitting is already minimal and reduced.

Analogously, for a tree Δ one can consider the full subgraph of the extension graph Δ^e, whose vertex set is the set of all elements of $\mathbb{G}(\Delta)$ which are conjugate to the canonical generators corresponding to vertices of Δ of degree more than 1 (which are exactly those which have non-abelian centralizers). Call this subgraph the *reduced extension graph* of Δ, and denote it by $\widetilde{\Delta}^e$.

The reduced extension graph will play an essential role in the classification of RAAGs up to commensurability. If $H < \mathbb{G}(\Delta)$ is a subgroup of finite index, then H intersects each cyclic subgroup associated to the vertex groups of the reduced extension graph in a non-trivial cyclic subgroup, that is $H \cap \langle a_i^g \rangle = \langle (a_i^{k_i})^g \rangle$ for some $k_i \in \mathbb{N}$, $g \in G$. On the other hand, by the description of centralisers in RAAGs, every element in H whose centraliser is non-abelian belongs to some cyclic subgroup $\langle (a_i^{k_i})^g \rangle = H \cap \langle a_i^g \rangle$. Since the set of elements with non-abelian centralisers is an invariant set up to isomorphism, it follows that the reduced extension graph is an algebraic invariant in the class of finite index subgroups of $\mathbb{G}(\Delta)$, that is if $K \simeq H$ and $H <_{fi} \mathbb{G}(\Delta)$, then the graph $T(K)$ whose vertex set is in one-to-one correspondence with maximal cyclic subgroups generated by elements of K with non-abelian centraliser and there is an edge (u, v) whenever the corresponding elements commute, is isomorphic to the reduced extension graph of $\mathbb{G}(\Delta)$ (see Lemma 3).

From this observation, one can deduce that many classes of RAAGs are not commensurable (without using the stronger fact that they are not quasi-isometric). For instance, RAAGs whose defining graphs are trees are not commensurable to RAAGs whose defining graph have cycles; or RAAGs defined by cycles of different lengths are not commensurable, etc. At this point, a couple of remarks are in order. Firstly, the above observation does not extend to the extension graph, that is, commensurable RAAGs may not have isomorphic extension graphs. For instance, we will show that the RAAGs defined by paths of length 3 and 4 are commensurable but their extension graphs are not isomorphic (leaves in the tree are also leaves in the extension graphs and the minimal distances in the extension graphs between leaves is 3 and 4 respectively). The reason here is that there are elements with abelian centralisers that are *not* powers of conjugates of the canonical generators so they are not accounted at all in the extension graph. Secondly, the isomorphism of the reduced extension graph is a necessary condition but it is by far not sufficient. All RAAGs defined by trees of diameter 4 have isomorphic reduced extension graphs but there are infinitely many non-commensurable classes.

In the next lemmas we notice that the two trees, the (reduced) extension graph and the Bass-Serre tree associated to the (reduced) centraliser splitting are equivalent and that the reduced extension graph and its quotient by the action of H is invariant up to isomorphism.

Lemma 1. *If Δ is a tree, then Δ^e is also a tree, which is isomorphic (as a graph) to the Bass-Serre tree T corresponding to the centralizer splitting of $\mathbb{G}(\Delta)$. Moreover, $\widetilde{\Delta}^e$ is a subtree of Δ^e, which is isomorphic (as a graph) to the Bass-Serre tree $\widetilde{T}$ corresponding to the reduced centralizer splitting of $\mathbb{G}(\Delta)$. The graph isomorphisms above are equivariant, in the sense that the action of $\mathbb{G}(\Delta)$ by conjugation on Δ^e (or $\widetilde{\Delta}^e$) corresponds to the natural action of $\mathbb{G}(\Delta)$ on the Bass-Serre tree of the centralizer splitting (reduced centralizer splitting, respectively).*

From now on we denote $G = \mathbb{G}(\Delta)$.

Proof. Every vertex of Δ^e has the form v^g, where v is some canonical generator of G, and $g \in G$. By Bass-Serre theory, vertices of T correspond to left cosets of centralizers of canonical generators of G. Define a morphism β from Δ^e to T by sending v^g to the vertex of the form $gC(v)$. This gives a bijection of the vertex sets. Two vertices $v_1^{g_1}$ and $v_2^{g_2}$ of Δ^e are connected by an edge iff $[v_1^{g_1}, v_2^{g_2}] = 1$, iff $[v_1, v_2] = 1$ and $v_1^{g_1} = v_1^g$, $v_2^{g_2} = v_2^g$ for some $g \in G$, iff $[v_1, v_2] = 1$ and $g_1^{-1}g \in C(v_1)$, $g_2^{-1}g \in C(v_2)$ for some $g \in G$, iff $[v_1, v_2] = 1$ and $g_1C(v_1) \cap g_2C(v_2) \neq \varnothing$. By Bass-Serre theory, since Δ is a tree and therefore there are no HNN-extensions appearing, two vertices $g_1C(v_1)$ and $g_2C(v_2)$ of T are connected by an edge iff $[v_1, v_2] = 1$ and $gC(v_1) = g_1C(v_1)$, $gC(v_2) = g_2C(v_2)$ for some $g \in G$, iff $[v_1, v_2] = 1$ and $g_1C(v_1) \cap g_2C(v_2) \neq \varnothing$. This shows that β is indeed a graph isomorphism. It is easy to see that it is equivariant, and that we can restrict β to an equivariant isomorphism between $\widetilde{\Delta}^e$ and $\widetilde{T}$. □

Suppose Δ is a finite tree, and K is a finite index subgroup of $\mathbb{G}(\Delta)$. Then $\mathbb{G}(\Delta)$ acts on the reduced extension graph $\widetilde{\Delta}^e$, and K acts on $\widetilde{\Delta}^e$ by restriction. Let $\Psi(K)$ be the quotient graph: $\Psi(K) = K\backslash\widetilde{\Delta}^e$. Note that there are natural projection graph morphisms

$$\gamma_K : \widetilde{\Delta}^e \to \Psi(K)$$

and

$$\delta_K : \Psi(K) \to \Delta.$$

It is easy to see that the graph $\Psi(K)$ is finite, since K has finite index in $\mathbb{G}(\Delta)$.

Below $C_K(w) = K \cap C_G(w)$ is defined even if $w \in G$ is not in K. In fact, since K has finite index in G, there exists a natural m such that $w^m \in K$, and then $C_K(w) = C_K(w^m)$.

Lemma 2. *Let v be a vertex of $\Psi(K)$, and w be any vertex of $\widetilde{\Delta}^e$ such that $\gamma_K(w) = v$ (we think of w as an element of G). Then*

1. *There is a bijection between the vertices of $\Psi(K)$ adjacent to v and the equivalence classes of vertices of $\widetilde{\Delta}^e$ adjacent to w, modulo conjugation by K.*
2. *There is a bijection between the edges of $\Psi(K)$ incident to v and the equivalence classes of vertices of $\widetilde{\Delta}^e$ adjacent to w, modulo conjugation by $C_K(w)$.*

Proof. The first claim follows immediately from the definition of $\Psi(K)$ as a quotient graph. Now let e_1 and e_2 be two different edges of $\widetilde{\Delta}^e$ incident to w. Let w_1, w_2 be the other ends of e_1, e_2 respectively. Then e_1 and e_2 project into the same edge of $\Psi(K)$ if and only if there exists $g \in K$ which takes e_1 to e_2. Since w and w_1 for sure belong to different orbits under the action of H (even of G), this happens if and only if g takes w_1 to w_2 and leaves w fixed. This means that $w_1^g = w_2$ and $w^g = w$, so $g \in C_K(w)$, thus the second claim also holds. □

In particular, graph $\Psi(K)$ can have multiple edges and cycles (but not loops).

Suppose now Δ and Δ' are finite trees, and denote $G = \mathbb{G}(\Delta)$ and $G' = \mathbb{G}(\Delta')$. Suppose G and G' are commensurable. Thus there exist finite index subgroups $H \leq G$ and $H' \leq G'$ and an isomorphism $\varphi : H \to H'$.

Consider the reduced extension graphs $\widetilde{\Delta}^e$ and $\widetilde{\Delta}'^e$ defined above, with the actions by conjugation of G and G' respectively. Finite graphs $\Psi(H)$ and $\Psi(H')$ are defined as above.

Lemma 3. *The group isomorphism $\varphi : H \to H'$ induces graph isomorphisms $\overline{\varphi} : \widetilde{\Delta}^e \to \widetilde{\Delta}'^e$ and $\varphi_* : \Psi(H) \to \Psi(H')$.*

Proof. Let $u_1, \dots, u_k$ be all vertices of Δ which have degree more than 1, and $u'_1, \dots, u'_{k'}$ be all vertices of Δ' which have degree more than 1. Then $u_1, \dots, u_k$ are all canonical generators of G with non-abelian centralizers, and $u'_1, \dots, u'_{k'}$ are all canonical generators of G' with non-abelian centralizers.

Notice that the only elements of G which have non-abelian centralizers in G are conjugates of powers of $u_1, \dots, u_k$ by some element of G. Since H has finite index in G and so $C_H(x) = H \cap C_G(x)$ has finite index in $C_G(x)$ for any $x \in G$, also the only elements of H which have non-abelian centralizers in H are conjugates of powers of $u_1, \dots, u_k$ by some element of G which belong to H. Analogously, the only elements of H' which have non-abelian centralizers in H' are conjugates of powers of $u'_1, \dots, u'_{k'}$ by some element of G' which belong to H'. Thus, the isomorphism φ should take the conjugates of powers of $u_1, \dots, u_k$ which are in H into the conjugates of powers of $u'_1, \dots, u'_{k'}$ which are in H'.

Now define a morphism of trees $\overline{\varphi} : \widetilde{\Delta}^e \to \widetilde{\Delta}'^e$ as follows. First define $\overline{\varphi}$ on the vertex set of $\widetilde{\Delta}^e$. Let w be a vertex of $\widetilde{\Delta}^e$. Then $w = u^g$ for some $g \in G$, where u is one of $u_1, \dots, u_k$. Since H has finite index in G, there exists a minimal positive integer k that $w^k \in H$. Then w^k has non-abelian centralizer in H, so $\varphi(w^k)$ also has non-abelian centralizer in H', thus $\varphi(w^k) = (u_0^l)^{g_0} \in H'$ for some positive integer l, $g_0 \in G'$, and u_0 equal to one of $u'_1, \dots, u'_{k'}$. Denote $w_0 = u_0^{g_0}$. Then let $\overline{\varphi}(w) = w_0$. Note that here l is also equal to the minimal positive integer i such that $w_0^i \in H'$, since if $i < l$, then also $i|l$, so w_0^l is a proper power in H', but this is impossible, since w^k is not a proper power in H, and $w_0^l = \varphi(w^k)$.

We can extend $\overline{\varphi}$ to the edges in the natural way. Indeed, two vertices w_1, w_2 of $\widetilde{\Delta}^e$ are adjacent iff $[w_1, w_2] = 1$, iff $[w_1^{k_1}, w_2^{k_2}] = 1$ (take positive integers k_1, k_2 such that $w_1^{k_1}, w_2^{k_2} \in H$), iff $[\varphi(w_1^{k_1}), \varphi(w_2^{k_2})] = 1$, iff $[\overline{\varphi}(w_1), \overline{\varphi}(w_2)] = 1$, iff $\overline{\varphi}(w_1)$ and $\overline{\varphi}(w_2)$ are adjacent. This shows that $\overline{\varphi}$ is a well-defined morphism of trees. Moreover, it is easy to see that $\overline{\varphi}$ is in fact isomorphism of trees, since the inverse morphism can be constructed in the same way.

If w is a vertex of $\widetilde{\Delta}^e$, and $h \in H$, then it is easy to see that we have $\overline{\varphi}(w^h) = \overline{\varphi}(w)^{\varphi(h)}$. It follows that we can restrict $\overline{\varphi}$ to the graph isomorphism

$$\varphi_* : \Psi(H) = H \backslash \widetilde{\Delta}^e \to \Psi(H') = H' \backslash \widetilde{\Delta}^e, \quad \varphi_*(\gamma_H(z)) = \gamma_{H'}(\overline{\varphi}(z)),$$

where z is a vertex or an edge of $\widetilde{\Delta}^e$, and γ_H, $\gamma_{H'}$ are the orbit projections as above. This proves the lemma. □

So far we have observed that isomorphisms leave the set of powers of conjugates of canonical generators with non-abelian centralisers invariant. However, if an isomorphism φ sends $(a^k)^g$ to $(a'^{k'})^{g'}$, then a priori there is no relation between the integer numbers k and k' corresponding to the powers.

A trivial example would be as follows. Let $H, K <_{fi} \mathbb{Z} \times F_2 \simeq \langle c \rangle \times \langle a, b \rangle$, $H = \langle c^k, a, b \rangle$ and $K = \langle c^{k'}, a, b \rangle$. Clearly, $\varphi : H \to K$ that maps $c^k \to c^{k'}$, $a \to a$ and $b \to b$ is an isomorphism and k and k' can be taken arbitrarily.

On the other hand, if we consider $H = \langle c^k, a^2, b, a^b\rangle$, $K = \langle c^{k'}\rangle \times F_m <_{fi} \langle c\rangle \times \langle a, b\rangle$ and we *assume* that the isomorphism $\varphi : H \to K$ sends c^k to $c^{k'}$ and each power of a conjugate of either a or b to a power of a conjugate of a and b, then we do get a constraint on the possible powers of the image of a. Indeed, the isomorphism φ induces an isomorphism φ' from the subgroup $\langle a^2, b, a^b\rangle$ to F_m, hence $m = 3$ and the index of $F_3 < F_2$ is 2. Since there are 3 conjugates of powers of generators in H, by our assumption, there are also 3 in K. Furthermore, since the sum of the powers of conjugates of a fixed generator is equal to the index, which in our case is 2, and since, upto relabeling, there are only two covers of F_2 of degree 2 and only one of them has 3 conjugates of powers of generators, it follows that either a is sent to a conjugate of a generator (and there are exactly two conjugates of this generator) or it is sent to a square of a generator (and it is the only conjugate of this generator in the subgroup).

Our next goal is to formalise and generalise these ideas to relate the powers of the different (conjugacy) classes of powers of generators that belong to the subgroup, with the indices of subgroups of centralisers in free groups. Since the isomorphism is going to send subgroups of a free group to subgroups of a free group, the rank of the subgroups is the same and using the relation with the corresponding indices, we will finally deduce a system of equations relating the powers of conjugates of generators that belong to the subgroups H and K.

Fix the subgroup $K \leq G$. We will now define labels of vertices and edges of $\widetilde{\Delta}^e$, and then of $\Psi(K)$.

Let w be a vertex of $\widetilde{\Delta}^e$, thus w is also an element of G. Define the *label of the vertex* w, denoted by $\overline{L}(w)$, to be the minimal positive integer k such that $w^k \in K$. Such number exists, since K has finite index in G.

For an edge f of $\widetilde{\Delta}^e$ connecting vertices w_1 and w_2 define the *label of the edge* f *at the vertex* w_1, denoted by $\bar{l}_{w_1}(f)$, to be the minimal positive integer k such that there exists an integer l such that $w_1^k w_2^l \in K$. Note that we can always suppose l is non-negative. Analogously the label of f at w_2 is defined. Note that by definition $\overline{L}(w_1) \geq \bar{l}_{w_1}(f)$, for all edges f.

Note that the labels of vertices and edges are invariant under the action of K on $\widetilde{\Delta}^e$ (by conjugation). Indeed, for $h \in K$ $w^k \in K$ iff $(w^h)^k \in K$, and $w_1^k w_2^l \in K$ iff $(w_1^h)^k (w_2^h)^l \in K$.

This means that we can define labels for the quotient graph $\Psi(K)$ as well. If v is a vertex of $\Psi(K)$, then define the *label of the vertex* v, denoted by $L(v)$, to be the label $\overline{L}(w)$, where w is some vertex of $\widetilde{\Delta}^e$ such that $\gamma_K(w) = v$. Analogously, if p is an edge of $\Psi(K)$ connecting vertices v_1 and v_2, then define the *label of the edge* p *at the vertex* v_1, denoted by $l_{v_1}(p)$, to be the label $\bar{l}_{w_1}(f)$, where f is some edge of $\widetilde{\Delta}^e$ such that $\gamma_K(f) = p$, and w_1 is the end of p such that $\gamma_K(w_1) = v_1$. These labels are correctly defined. Note that the labels of vertices and edges of $\Psi(K)$ are positive integers.

Lemma 4. *Let p be an edge of $\Psi(K)$ connecting vertices v_1 and v_2. Then*

$$\frac{L(v_1)}{l_{v_1}(p)} = \frac{L(v_2)}{l_{v_2}(p)} = r, \tag{1}$$

where r is some positive integer.

Proof. Let f be some edge of $\widetilde{\Delta}^e$ such that $\gamma_K(f) = p$, and w_1, w_2 be the ends of f, so that $\gamma_K(w_1) = v_1$, $\gamma_K(w_2) = v_2$. Then we can rewrite (1) as

$$\frac{\overline{L}(w_1)}{\overline{l}_{w_1}(f)} = \frac{\overline{L}(w_2)}{\overline{l}_{w_2}(f)} = r. \tag{2}$$

By definition $\overline{l}_{w_1}(f)$ is the minimal positive integer k such that there exists an integer l such that $w_1^k w_2^l \in K$. Note that if k' is some other positive integer such that there exists an integer l' such that $w_1^{k'} w_2^{l'} \in K$, then $\overline{l}_{w_1}(f)|k'$. Indeed, denote $k_0 = \overline{l}_{w_1}(f)$ and let d be the greatest common divisor of k_0 and k'. Then there exist integers α, β such that $\alpha k_0 + \beta k' = d$. We have $w_1^{k_0} w_2^l \in K$ and $w_1^{k'} w_2^{l'} \in K$, so, since w_1 and w_2 commute,

$$(w_1^{k_0} w_2^l)^\alpha (w_1^{k'} w_2^{l'})^\beta = w_1^{\alpha k_0 + \beta k'} w_2^{\alpha l + \beta l'} = w_1^d w_2^{\alpha l + \beta l'} \in K,$$

so $d \geq k_0$, but since $d = gcd(k_0, k')$, this means that $d = k_0$ and thus $k_0 = \overline{l}_{w_1}(f)|k'$.

Recall that $\overline{L}(w_1)$ is the minimal positive integer k such that $w_1^k \in K$, so $w_1^{\overline{L}(w_1)} \in K$ and it follows from above that $\overline{l}_{w_1}(f)|\overline{L}(w_1)$. Thus, $\frac{\overline{L}(w_1)}{\overline{l}_{w_1}(f)} = r_1$ is a positive integer. Analogously $\frac{\overline{L}(w_2)}{\overline{l}_{w_2}(f)} = r_2$ is a positive integer.

It remains to show that $r_1 = r_2$. Indeed, suppose that l is such that $w_1^{\overline{l}_{w_1}(f)} w_2^l \in K$. As shown above, this means that $\overline{l}_{w_2}(f)|l$, so let $l = \overline{l}_{w_2}(f)q$. We have

$$(w_1^{\overline{l}_{w_1}(f)} w_2^l)^{r_2} = w_1^{\overline{l}_{w_1}(f) r_2} w_2^{l r_2} = w_1^{\overline{l}_{w_1}(f) r_2} w_2^{\overline{l}_{w_2}(f) q r_2} = w_1^{\overline{l}_{w_1}(f) r_2} w_2^{\overline{L}(w_2) q} \in K.$$

But $w_2^{\overline{L}(w_2)} \in K$, so $w_2^{\overline{L}(w_2) q} \in K$. Thus $w_1^{\overline{l}_{w_1}(f) r_2} \in K$. This means that $\overline{l}_{w_1}(f) r_2 \geq \overline{L}(w_1) = \overline{l}_{w_1}(f) r_1$, so $r_2 \geq r_1$. Analogous argument shows that $r_1 \geq r_2$. Thus $r_1 = r_2$. □

Let w be a vertex of $\widetilde{\Delta}^e$, and let $\gamma_K(w) = v$ and $u = \delta_K(v)$. Thus u is a canonical generator of G with non-abelian centralizer, and $w = u^g$ for some $g \in G$. Suppose $u_1, \dots, u_k$ are all vertices of Δ adjacent to u, then $k \geq 2$ is the degree of u in Δ, and $u_1, \dots, u_s$ are those of them which have degree more than 1 (and so have non-abelian centralizer), here $s \leq k$. Note that

$$C_K(w) \leq C_G(w) \cong Z(C_G(w)) \times F(u_1^g, \dots, u_k^g),$$

where $Z(C_G(w)) = \langle w \rangle$ is cyclic and $F(u_1^g, \dots, u_k^g)$ is the free group of rank k with the basis $u_1^g, \dots, u_k^g$. Let $F_{K,w}$ be the free group of rank k with the basis $q_1, \dots, q_k$. Denote by $\pi : C_G(w) \to F_{K,w}$ the factorization by the center homomorphism: $\pi(w) = 1$, $\pi(u_i^g) = q_i$, $i = 1, \dots, k$. Then π induces an isomorphism between $F(u_1^g, \dots, u_k^g)$ and $F_{K,w}$. Below, when we speak about cycles in the Schreier graph, we mean simple cycles with all edges labelled by the same generator, and we mean the Schreier graph with respect to the generators $q_1, \dots, q_k$ of $F_{K,w}$.

Lemma 5. *In the above notations the following statements hold:*

1. *π induces a group embedding $\pi_C : C_K(w)/Z(C_K(w)) \hookrightarrow F_{K,w}$,*
   and $P_{K,w} = \pi_C(C_K(w)/Z(C_K(w))) = \pi(C_K(w))$ is a finite index subgroup of $F_{K,w}$.
2. *π induces a bijection π_* between the edges of $\Psi(K)$ incident to $v = \gamma_K(w)$ and those cycles in the Schreier graph of $P_{K,w}$ in $F_{K,w}$ which correspond to the generators $q_1, \dots, q_s$.*

(3) *For every edge p of $\Psi(K)$ connecting $v = \gamma_K(w)$ with some other vertex v_1, the label $l_{v_1}(p)$ of p with respect to v_1 is equal to the length of the cycle $\pi_*(p)$ in the Schreier graph of $P_{K,w}$ in $F_{K,w}$.*

(4) *If w, w' are two vertices of $\widetilde{\Delta}^e$ in the same orbit under the action of K, then there is a natural isomorphism α between $F_{K,w}$ and $F_{K,w'}$, which restricts to an isomorphism α_P between $P_{K,w}$ and $P_{K,w'}$ and respects the bijections from above.*

Proof. Let j be the minimal (positive) power of w which belongs to K. Note that $Z(C_K(w)) \leq Z(C_G(w))$, since if some $z \in C_G(w)$, then for some positive integer l $z^l \in C_K(w)$, so every element in the center of $C_K(w)$ should also commute with z. Also $Ker(\pi) = Z(C_G(w)) = \langle w \rangle$, so $Ker(\pi|_{C_K(w)}) = K \cap Z(C_G(w)) = \langle w^j \rangle = Z(C_K(w))$. This means that π induces an embedding $\pi_C : C_K(w)/Z(C_K(w)) \hookrightarrow F_{K,w}$. Since $C_K(w) = K \cap C_G(w)$ has finite index in $C_G(w)$, it follows that $P_{K,w}$ has finite index in $F_{K,w}$. This proves the first statement.

Now we define a bijection $\overline{\pi}$ between the edges of $\widetilde{\Delta}^e$ incident to w and lines in the Cayley graph of $F_{K,w}$ (with respect to the basis $q_1, \dots, q_k$) consisting of edges labelled by the same generator, one of $q_1, \dots, q_s$ (which have non-abelian centralizers). Let x be one of the generators $u_1, \dots, u_s$, and f be an edge connecting $w = u^g$ with $w_1 = x^{g_1}$, for some $g_1 \in G$. As in the proof of Lemma 1, this means that $gC(u) \cap g_1 C(x) \neq \varnothing$, so $w_1 = x^{g_1} = x^{gg_2}$, where $g_2 \in C(u) = \langle u, u_1, \dots, u_k \rangle$. Note that g_2 is defined up to multiplication on the right by a power of x. We can suppose that $g_2 \in \langle u_1, \dots, u_k \rangle$, since u commutes with $u_1, \dots, u_k$, in particular with x as well. Then $w_1 = gg_2 x g_2^{-1} g^{-1} = yx^g y^{-1}$, where $y = g_2^g$ is the word obtained from g_2 by replacing each u_i by u_i^g, for $i = 1, \dots, k$, so y can be thought of as an element of $F(u_1^g, \dots, u_k^g)$. Now let $\overline{\pi}(f)$ be the line in the Cayley graph of $F_{K,w}$ passing through the vertex corresponding to $\pi(y)$, and with edges labeled by the element $\pi(x^g) = q$, which is equal to one of the elements $q_1, \dots, q_s$.

Note that $\overline{\pi}$ is well-defined. Indeed, suppose we can write w_1 in two ways: $w_1 = x^{g_1} = x^{gg_2}$ and $w_1 = x^{g_1'} = x^{gg_2'}$, where $g_2, g_2' \in \langle u_1, \dots, u_k \rangle$, and x is one of $u_1, \dots, u_s$. Thus in G we have $x^{gg_2} = x^{gg_2'}$, so $x^{g_2} = x^{g_2'}$. This can be considered as an equality in the free group on $u_1, \dots, u_k$. Then $[x, g_2^{-1} g_2'] = 1$, so $g_2^{-1} g_2' = x^k$ for some $k \in \mathbb{Z}$, which means that $g_2' = g_2 x^k$, and then $y' = y(x^g)^k$, where $y = g_2^g$, $y' = g_2'^g$, which means that $\pi(y') = \pi(y) q^k$, so the line in the Cayley graph is defined correctly.

It is easy to see from the definition of $\overline{\pi}$ that it is surjective. Now show that $\overline{\pi}$ is injective. Suppose f connects w with $w_1 = x^{g_1}$, anf f' connects w with $w_1' = x^{g_1'}$, $g_1 = gg_2$, $y = g_2^g$ and $g_1' = gg_2'$, $y' = g_2'^g$, where $g_2, g_2' \in \langle u_1, \dots, u_k \rangle$. If $\overline{\pi}(f) = \overline{\pi}(f')$, then $\pi(y) = \pi(y') q^k$, so $y = y'(x^g)^k$ for some $k \in \mathbb{Z}$. Then $g_2 = g_2' x^k$, and thus $g_1 = g_1' x^k$, so $w_1 = w_1'$. This shows that $\overline{\pi}$ is bijective.

Note that $C_G(w)$ acts on the set of edges incident to the vertex w of $\widetilde{\Delta}^e$, and thus $C_K(w)$ also acts on this set. The action of w is trivial, so this induces the action of $C_K(w)/Z(C_K(w))$ on the set of edges incident to w. On the other hand, $P_{K,w} = \pi_C(C_K(w)/Z(C_K(w))) \leq F_{K,w}$ acts on the Cayley graph of $F_{K,w}$ in the natural way (by left multiplication), and so $P_{K,w}$ also acts on the set of lines in the Cayley graph of $F_{K,w}$ defined above. Recall that π_C induces an isomorphism $C_K(w)/Z(C_K(w)) \cong \pi_C(C_K(w)/Z(C_K(w)))$. It is easy to see from the definition of $\overline{\pi}$ above that for every edge f of $\widetilde{\Delta}^e$ incident to w and every $h \in C_K(w)/Z(C_K(w))$ we have $\overline{\pi}(h \cdot f) = \pi_C(h) \cdot \overline{\pi}(f)$, where we denote both actions defined above by dots. This means that the bijection $\overline{\pi}$ induces a

bijection between the orbits of the edges incident to w under the action of $C_K(w)$ on one side, and the orbits of the lines (as above) in the Cayley graph of $F_{K,w}$ under the action of $P_{K,w}$ on the other side. By Lemma 2, the orbits of the edges incident to w under the action of $C_K(w)$ are in one-to-one correspondence with the edges of $\Psi(K)$ incident to $v = \gamma_K(w)$. And the orbits of the lines (as above) in the Cayley graph of $F_{K,w}$ under the action of $P_{K,w}$ are in a natural one-to-one correspondence with the cycles in the Schreier graph of $P_{K,w}$ in $F_{K,w}$, which are labelled by the generators $q_1, \dots, q_s$. This defines the desired bijection π_* and proves the second claim of the Lemma.

Now we turn to the third claim. Let f be an edge of $\widetilde{\Delta}^e$ incident to w such that $\gamma_K(f) = p$, and let f connect w with w_1, so $\gamma_K(w_1) = v_1$. Recall that the label of p at v_1 is equal to the label of f at w_1, $\bar{l}_{w_1}(f)$, which is the minimal positive integer k such that there exists an integer l such that $w_1^k w^l \in K$. Note that this means that $\pi(w_1^k) = \pi(w_1^k w^l) \in P_{K,w}$. Since $Ker(\pi) = \langle w \rangle$, it follows that $\bar{l}_{w_1}(f)$ is equal to the minimal positive integer k' such that $(\pi(w_1))^{k'} \in P_{K,w}$. Let $w = u^g$ and $w_1 = x^{gg_2}$ as above, where $g_2 \in \langle u_1, \dots, u_k \rangle$.

Denote, as above, $\pi(x^g) = q$, $g_2^g = y$. We have $\pi(w_1) = \pi(x^{gg_2}) = \pi(y)q\pi(y)^{-1}$, and $\overline{\pi}(f)$ is the line passing through $\pi(y)$ and with edges labelled by q. Thus $\bar{l}_{w_1}(f)$ is equal to the minimal positive integer k' such that

$$(\pi(w_1))^{k'} = (\pi(y)q\pi(y)^{-1})^{k'} = \pi(y)q^{k'}\pi(y)^{-1} \in P_{K,w},$$

which is equivalent to $r\pi(y) = \pi(y)q^{k'}$ for some $r \in P_{K,w}$. Thus, $\bar{l}_{w_1}(f)$ is equal to the minimal positive exponent k' of q for which points $\pi(y)$ and $\pi(y)q^{k'}$ on the line $\overline{\pi}(f)$ in the Cayley graph of $F_{K,w}$ are equivalent under the action of $P_{K,w}$, and this is exactly the length of the corresponding cycle $\pi_*(f)$ in the Schreier graph of $P_{K,w}$ in $F_{K,w}$. This proves the third claim.

It remains to prove the last claim. Let $w' = w^h$, where $h \in K$, and let $u, u_1, \dots, u_k$ be as above. Then

$$w = u^g,\ C_G(w) \cong \langle u^g \rangle \times F(u_1^g, \dots, u_k^g),$$
$$w' = u^{hg},\ C_G(w') = (C_G(w))^h \cong \langle u^{hg} \rangle \times F(u_1^{hg}, \dots, u_k^{hg}).$$

Also

$$F_{K,w} = F(q_1, \dots, q_k),\ F_{K,w'} = F(q_1', \dots, q_k'),$$
$$\pi : C_G(w) \to F_{K,w},\ \pi(w) = 1,\ \pi(u_i^g) = q_i,\ i = 1, \dots, k,$$
$$\pi' : C_G(w') \to F_{K,w'},\ \pi(w^h) = 1,\ \pi(u_i^{hg}) = q_i',\ i = 1, \dots, k.$$

Let $\overline{\alpha}$ be the isomorphism $\overline{\alpha} : C_G(w) \to C_G(w')$, $\overline{\alpha}(t) = t^h$ for $t \in C_G(w)$. Define the isomorphism $\alpha : F_{K,w} \to F_{K,w'}$, $\alpha(q_i) = q_i'$, $i = 1, \dots, k$. Then $\alpha\pi = \pi'\overline{\alpha}$ by definitions.

We have

$$C_K(w') = K \cap C_G(w') = K \cap (C_G(w))^h = (K \cap C_G(w))^h = C_K(w)^h,$$

so $\overline{\alpha}(C_K(w)) = C_K(w')$. Note that $P_{K,w} = \pi(C_K(w))$, $P_{K,w'} = \pi'(C_K(w'))$, so

$$\alpha(P_{K,w}) = \alpha\pi(C_K(w)) = \pi'\overline{\alpha}(C_K(w)) = \pi'(C_K(w')) = P_{K,w'},$$

so α induces an isomorphism α_P between $P_{K,w}$ and $P_{K,w'}$. Thus α also induces an isomorphism between the Schreier graphs of $P_{K,w}$ in $F_{K,w}$ and of $P_{K,w'}$ in $F_{K,w'}$ in a natural way.

Recall that π_* is a bijection between the edges of $\Psi(K)$ incident to $v = \gamma_K(w)$ and those cycles in the Schreier graph of $P_{K,w}$ in $F_{K,w}$ which correspond to the generators $q_1, \dots, q_s$. Analogously π'_*

is a bijection between the edges of $\Psi(K)$ incident to $v = \gamma_K(w') = \gamma_K(w)$ and those cycles in the Schreier graph of $P_{K,w'}$ in $F_{K,w'}$ which correspond to the generators $q'_1, \dots, q'_s$. It is easy to see from the definition of π_*, π'_* that for every edge p of $\Psi(K)$ incident to v the cycle $\pi_*(p)$ goes into the cycle $\pi'_*(p)$ under the isomorphism induced by α, i.e., α respects the bijections. □

If v is some vertex of $\Psi(H)$, then by the last claim of Lemma 5, a different choice of w in $\widetilde{\Delta}^e$ such that $\gamma_K(w) = v$ gives the same groups $F_{K,w}$, subgroups $P_{K,w}$, as well as bijections π_*, as in Lemma 5, up to isomorphism. This means that the exact choice of such w is inessential, and, abusing the notation, we will denote $F_{K,v} = F_{K,w}$, $P_{K,v} = P_{K,w}$, where the vertex w is any vertex of $\widetilde{\Delta}^e$ such that $\gamma_K(w) = v$.

Lemma 6. *In the above notation, with v being a vertex of $\Psi(K)$ and $u = \delta_K(v)$, let u_0 be some vertex of Δ of degree more than 1, connected to u by an edge e_0. If $f_1, \dots, f_k$ are all edges of $\Psi(K)$ incident to the vertex v such that $\delta_K(f_i) = e_0$ for all $i = 1, \dots, k$, and each f_i connects v with the vertex v_i $(i = 1, \dots, k)$, then*

$$\sum_{i=1}^{k} l_{v_i}(f_i) = |F_{K,v} : P_{K,v}|. \tag{3}$$

If $z_1, \dots, z_p$ are all edges of $\Psi(K)$ incident to v, and each z_i connects v with the vertex w_i, then

$$\sum_{i=1}^{p} l_{w_i}(z_i) = m\,|F_{K,v} : P_{K,v}|. \tag{4}$$

where m is the number of vertices of degree greater than 1 adjacent to u in Δ.

Proof. According to Lemma 5, there is a bijection between the set of edges $f_1, \dots, f_k$ and the set of cycles in the Schreier graph of $P_{K,v}$ in $F_{K,v}$ which correspond to conjugates of u_0, and the length of each cycle is equal to the corresponding edge label. The sum of lengths of these cycles is equal to the number of vertices in the Schreier graph, which is equal to $|F_{K,v} : P_{K,v}|$, thus (3) follows. Summing all the equalities of the form (3) for all vertices of Δ of degree greater than 1 adjacent to u gives (4). □

Lemma 7. *Let v be a vertex of $\Psi(H)$, and let $v' = \varphi_*(v)$ be the corresponding vertex of $\Psi(H')$. Let $\delta_H(v) = u$ and $\delta_{H'}(v') = u'$, and suppose u has degree t in Δ, and u' has degree t' in Δ'. Then, in the notation of* Lemmas 5 *and* 6*,*

$$(t-1)\,|F_{H,v} : P_{H,v}| = (t'-1)\,|F_{H',v'} : P_{H',v'}|. \tag{5}$$

Proof. Let $w \in \widetilde{\Delta}^e$ be such that $\gamma_H(w) = v$, and think of w as an element of G. Let l be such positive integer that $w^l \in H$. Then

$$C_H(w) = C_H(w^l) \cong C_{H'}(\varphi(w^l)) = C_{H'}(\varphi(w)).$$

Note that $\gamma_{H'}(\varphi(w)) = v'$. Thus

$$P_{H,v} \cong C_H(w)/Z(C_H(w)) \cong C_{H'}(\varphi(w))/Z(C_{H'}(\varphi(w))) \cong P_{H',v'}.$$

In particular, $rk(P_{H,v}) = rk(P_{H',v'})$. Then, by Schreier index formula,

$$|F_{H,v} : P_{H,v}|(rk(F_{H,v}) - 1) = rk(P_{H,v}) - 1 = rk(P_{H',v'}) - 1 = |F_{H',v'} : P_{H',v'}|(rk(F_{H',v'}) - 1).$$

This implies (5). □

Let f be an edge of $\widetilde{\Delta}^e$ incident to a vertex w, and $f' = \overline{\varphi}(f)$, $w' = \overline{\varphi}(w)$. By $\overline{L}(w)$, $\overline{L'}(w)$ we denote the labels of the vertices w in $\widetilde{\Delta}^e$, w' in $\widetilde{\Delta}'^e$ respectively. By $\bar{l}_w(f)$ we denote the label of the edge f with respect to w in $\widetilde{\Delta}^e$, and by $\overline{l'}_w(f)$ we denote the label of the edge f' with respect to w' in $\widetilde{\Delta}'^e$.

Analogously, let p be an edge of $\Psi(H)$ incident to a vertex v, and $p' = \varphi_*(p)$, $v' = \varphi_*(v)$. By $L(v)$, $L'(v)$ we denote the labels of the vertices v in $\Psi(H)$, v' in $\Psi(H')$ respectively. By $l_v(p)$ we denote the label of the edge p with respect to v in $\Psi(H)$, and by $l'_v(p)$ we denote the label of the edge p' with respect to v' in $\Psi(H')$.

Lemma 8. *Let p be an edge of $\Psi(H)$ connecting vertices v_1 and v_2. Then*

$$\frac{L(v_1)}{l_{v_1}(p)} = \frac{L'(v_1)}{l'_{v_1}(p)} = \frac{L(v_2)}{l_{v_2}(p)} = \frac{L'(v_2)}{l'_{v_2}(p)} = q, \tag{6}$$

where q is some positive integer.

Proof. We already know from Lemma 4 that

$$\frac{L(v_1)}{l_{v_1}(p)} = \frac{L(v_2)}{l_{v_2}(p)} = q_1, \quad \frac{L'(v_1)}{l'_{v_1}(p)} = \frac{L'(v_2)}{l'_{v_2}(p)} = q_2,$$

where q_1 and q_2 are some positive integers. Thus it suffices to show that $q_1 = q_2$, or

$$\frac{L(v_1)}{l_{v_1}(p)} = \frac{L'(v_1)}{l'_{v_1}(p)}. \tag{7}$$

Let f be some edge in $\widetilde{\Delta}^e$ such that $\gamma_H(f) = p$, and w_1, w_2 be the ends of f, so that $\gamma_H(w_1) = v_1$, $\gamma_H(w_2) = v_2$. Let also $w'_1 = \overline{\varphi}(w_1)$, $w'_2 = \overline{\varphi}(w_2)$. Then we can rewrite (7) as

$$\frac{\overline{L}(w_1)}{\bar{l}_{w_1}(f)} = \frac{\overline{L'}(w_1)}{\overline{l'}_{w_1}(f)}. \tag{8}$$

Recall that $\bar{l}_{w_1}(f)$ is the minimal positive integer k such that there exists an integer l such that $w_1^k w_2^l \in H$. Fix such number l. Denote $\overline{L}(w_1) = m_1$, $\overline{L}(w_2) = m_2$, $\overline{L'}(w_1) = m'_1$, $\overline{L'}(w_2) = m'_2$. Then $w_1^{\bar{l}_{w_1}(f)} w_2^l \in H$, and the element $y = (w_1^{\bar{l}_{w_1}(f)} w_2^l)^{m_1 m_2}$ is a $m_1 m_2$ power in H. Note that

$$y = (w_1^{m_1})^{\bar{l}_{w_1}(f) m_2} \, (w_2^{m_2})^{l m_1}.$$

Note also that $w_1^{m_1}, w_2^{m_2} \in H$ by definition of labels, and $\varphi(w_1^{m_1}) = (w'_1)^{m'_1}$, $\varphi(w_2^{m_2}) = (w'_2)^{m'_2}$, as shown in the proof of Lemma 3. Thus

$$\varphi(y) = ((w'_1)^{m'_1})^{\bar{l}_{w_1}(f) m_2} \, ((w'_2)^{m'_2})^{l m_1}$$

is a $m_1 m_2$ power in H': $\varphi(y) = z^{m_1 m_2}$ for some $z \in H'$. But roots in RAAGs are unique, see [DK93], so

$$z = (w'_1)^{m'_1 \bar{l}_{w_1}(f)/m_1} \, (w'_2)^{m'_2 l/m_2} \in H'.$$

By definition of $\overline{l'}_{w_1}(f)$, this means that $m'_1 \bar{l}_{w_1}(f)/m_1 \geq \overline{l'}_{w_1}(f)$, so

$$\frac{\overline{L'}(w_1)}{\overline{l'}_{w_1}(f)} \geq \frac{\overline{L}(w_1)}{\bar{l}_{w_1}(f)}.$$

The same argument with the roles of G and G' interchanged shows that

$$\frac{\overline{L}(w_1)}{\overline{l}_{w_1}(f)} \geq \frac{\overline{L'}(w_1)}{\overline{l'}_{w_1}(f)}.$$

Thus

$$\frac{\overline{L}(w_1)}{\overline{l}_{w_1}(f)} = \frac{\overline{L'}(w_1)}{\overline{l'}_{w_1}(f)}.$$

This proves the Lemma. □

The above lemmas imply that the labels of edges satisfy the following equations.

Lemma 9. *Suppose v is some vertex of $\Psi(H)$, and $v' = \varphi_*(v)$. Suppose $\delta_H(v)$ has degree t in Δ and is adjacent to r vertices of degree more than 1 in Δ, and $\delta_{H'}(v')$ has degree t' in Δ', and is adjacent to r' vertices of degree more than 1 in Δ'. Let $f_1, \dots, f_k$ be all edges of $\Psi(H)$ beginning in v, and suppose $\alpha_i = l_v(f_i)$, $\alpha'_i = l'_v(f_i)$, $i = 1, \dots, k$, are their labels. Then we have*

$$\frac{t-1}{r} \sum_{i=1}^{k} \alpha_i L(v_i) = \frac{t'-1}{r'} \sum_{i=1}^{k} \alpha_i L'(v_i), \tag{9}$$

$$\frac{t-1}{r} \sum_{i=1}^{k} \alpha'_i L(v_i) = \frac{t'-1}{r'} \sum_{i=1}^{k} \alpha'_i L'(v_i). \tag{10}$$

Proof. Let the edge f_i connect v to v_i, $i = 1, \dots, k$. By Lemma 6 we have

$$\sum_{i=1}^{k} l_{v_i}(f_i) = r|F_{H,v} : P_{H,v}|, \quad \sum_{i=1}^{k} l'_{v_i}(f_i) = r'|F_{H',v'} : P_{H',v'}|,$$

and from Lemma 7 we get

$$\frac{t-1}{r} \sum_{i=1}^{k} l_{v_i}(f_i) = \frac{t'-1}{r'} \sum_{i=1}^{k} l'_{v_i}(f_i).$$

Multiplying both sides by $L(v)$, we obtain

$$\frac{t-1}{r} \sum_{i=1}^{k} l_{v_i}(f_i)L(v) = \frac{t'-1}{r'} \sum_{i=1}^{k} l'_{v_i}(f_i)L(v). \tag{11}$$

Multiplying both sides by $L'(v)$, we obtain

$$\frac{t-1}{r} \sum_{i=1}^{k} l_{v_i}(f_i)L'(v) = \frac{t'-1}{r'} \sum_{i=1}^{k} l'_{v_i}(f_i)L'(v). \tag{12}$$

According to Lemma 8, we have

$$\frac{L(v)}{l_v(f_i)} = \frac{L'(v)}{l'_v(f_i)} = \frac{L(v_i)}{l_{v_i}(f_i)} = \frac{L'(v_i)}{l'_{v_i}(f_i)}$$

for all $i = 1, \dots k$, so

$$l_{v_i}(f_i)L(v) = L(v_i)l_v(f_i), \quad l'_{v_i}(f_i)L(v) = L'(v_i)l_v(f_i),$$
$$l_{v_i}(f_i)L'(v) = L(v_i)l'_v(f_i), \quad l'_{v_i}(f_i)L'(v) = L'(v_i)l'_v(f_i).$$

Thus (11), (12) imply (9), (10). □

If v is a vertex of $\Psi(H)$ such that $\delta_H(v) = u_1$, and $\delta_{H'}(\varphi_*(v)) = u_2$, then we say that v *has type* u_1/u_2, or v *is a* u_1/u_2 *vertex*.

We now turn our attention to the proofs of the main results. We first show that there are infinitely many different commensurability classes inside the class of RAAGs defined by trees of diameter 4. This is a particular case of Theorem 4 , but we give a separate proof for two reasons: the first one, it is technically easier and it helps as a warm up for the general case; secondly, for the reader interested only in the quantitative aspect of the result, it suffices to read our Theorem 3.

For $0 < d_1 \leq d_2$ we denote by P_{d_1,d_2} the tree of diameter 4 with 2 pivots, one of degree d_1 and the other of degree d_2, and no hair vertices, see Figure 2. Thus $P_{d_1,d_2} = T((d_1,1),(d_2,1);0)$ if $d_1 < d_2$ and $P_{d_1,d_1} = T((d_1,2);0)$.



FIGURE 2. Tree of diameter 4 with 2 pivots

Theorem 3 (Infinitely many commensurability classes)**.** *Let* $0 < m_1 \leq m_2$ *and* $0 < n_1 \leq n_2$. *Suppose the groups* $G = \mathbb{G}(P_{m_1,m_2})$ *and* $G' = \mathbb{G}(P_{n_1,n_2})$ *are commensurable. Then* $\frac{m_1}{n_1} = \frac{m_2}{n_2}$.

In particular, there are infinitely many commensurability classes among RAAGs defined by trees of diameter 4.

In fact, we will see in the next section that $G = \mathbb{G}(P_{m_1,m_2})$ and $G' = \mathbb{G}(P_{n_1,n_2})$ are commensurable if and only if $\frac{m_1}{n_1} = \frac{m_2}{n_2}$.

Proof of Theorem 3. Let c be the center vertex of P_{m_1,m_2} and c' be the center vertex of P_{n_1,n_2}. Let p_1, p_2 be the pivots of P_{m_1,m_2} of degrees m_1+1, m_2+1 correspondingly, and p_1', p_2' be the pivots of P_{n_1,n_2} of degrees n_1+1, n_2+1 correspondingly.

Let $H \leq G$ and $H' \leq G'$ be finite index subgroups such that $H \cong H'$. From Lemma 9 and Lemma 6 applied to both $\Delta = P_{m_1,m_2}$ and $\Delta' = P_{n_1,n_2}$ we have a system of equations on the labels of vertices and edges of $\Psi(H)$. We will show that this system of equations does not have any positive integer solutions unless $\frac{m_1}{n_1} = \frac{m_2}{n_2}$.

Note that there could exist the following 9 types of vertices in $\Psi(H)$: p_1/p_1', p_1/p_2', p_2/p_1', p_2/p_2', c/p_1', c/p_2', p_1/c', p_2/c', c/c'. It is easy to see that the following statements hold:

- Every p_1/p_1', p_1/p_2', p_2/p_1', p_2/p_2' vertex can be connected only with c/c' vertices;
- Every c/c' vertex can be connected only with p_1/p_1', p_1/p_2', p_2/p_1', p_2/p_2' vertices;
- Every p_1/c' and p_2/c' vertex can be connected only with c/p_1' and c/p_2' vertices;
- Every c/p_1' and c/p_2' vertex can be connected only with p_1/c' and p_2/c' vertices.

Thus, there are two cases: either $\Psi(H)$ has only p_1/p_1', p_1/p_2', p_2/p_1', p_2/p_2' and c/c' vertices, or $\Psi(H)$ has only p_1/c', p_2/c', c/p_1' and c/p_2' vertices.

Case 1. Suppose first $\Psi(H)$ has only p_1/p_1', p_1/p_2', p_2/p_1', p_2/p_2' and c/c' vertices.

Let v be a p_i/p_j' vertex, where $i = 1, 2$, $j = 1, 2$. Then all vertices adjacent to v have type c/c'. Let $f_1, \dots f_S$ be all edges incident to v, and let f_s connect v with v_s for $s = 1, \dots, S$ (possibly some of v_i coincide). Then, by Lemma 9, in the notations of which we have $t = m_i+1, r = 1, t' = n_j+1, r' = 1$, we have

$$m_i \sum_{s=1}^{S} L(v_s) l_v(f_s) = n_j \sum_{s=1}^{S} L'(v_s) l_v(f_s). \tag{13}$$

Now suppose that $w_1, \dots, w_Q$ are all c/c' vertices of $\Psi(H)$. For every vertex w_q, $q = 1, \dots, Q$, consider all the edges incident to w_q which finish in p_i/p_j' vertices (for fixed $i = 1, 2$ and $j = 1, 2$), and let X_{ij}^q be the sum of the l-labels of these edges at the p_i/p_j' ends; we let $X_{ij}^q = 0$ if there are no such edges. It follows from Lemma 6 that for all $q = 1, \dots, Q$ we have

$$X_{11}^q + X_{12}^q = |F_{H,w_j} : P_{H,w_j}| = X_{21}^q + X_{22}^q. \tag{14}$$

Now sum up the equalities of the form (13) for all p_i/p_j' vertices v (for fixed $i = 1, 2$ and $j = 1, 2$), and group the summands with the vertex label corresponding to the same c/c' vertex together. We obtain

$$m_1 \sum_{q=1}^{Q} L(w_q) X_{11}^q = n_1 \sum_{q=1}^{Q} L'(w_q) X_{11}^q, \tag{15}$$

$$m_1 \sum_{q=1}^{Q} L(w_q) X_{12}^q = n_2 \sum_{q=1}^{Q} L'(w_q) X_{12}^q, \tag{16}$$

$$m_2 \sum_{q=1}^{Q} L(w_q) X_{21}^q = n_1 \sum_{q=1}^{Q} L'(w_q) X_{21}^q, \tag{17}$$

$$m_2 \sum_{q=1}^{Q} L(w_q) X_{22}^q = n_2 \sum_{q=1}^{Q} L'(w_q) X_{22}^q. \tag{18}$$

Summing (15) and (16) and using that $n_1 \le n_2$, we get

$$m_1 \sum_{q=1}^{Q} L(w_j)(X_{11}^q + X_{12}^q) = n_1 \sum_{q=1}^{Q} L'(w_q) X_{11}^q + n_2 \sum_{q=1}^{Q} L'(w_q) X_{12}^q \ge n_1 \sum_{q=1}^{Q} L'(w_q)(X_{11}^q + X_{12}^q). \tag{19}$$

Analogously, summing (18) and (17) and using that $n_1 \leq n_2$, we get

$$m_2 \sum_{q=1}^{Q} L(w_q)(X_{21}^q + X_{22}^q) = n_1 \sum_{q=1}^{Q} L'(w_q) X_{21}^q + n_2 \sum_{q=1}^{Q} L'(w_q) X_{22}^q \leq n_2 \sum_{q=1}^{Q} L'(w_q)(X_{21}^q + X_{22}^q). \tag{20}$$

Multiplying (19) by m_2 and (20) by m_1 and using (14), we obtain

$$m_2 n_1 \sum_{q=1}^{Q} L'(w_q)(X_{11}^q + X_{12}^q) \leq m_2 m_1 \sum_{q=1}^{Q} L(w_q)(X_{11}^q + X_{12}^q) =$$

$$= m_1 m_2 \sum_{q=1}^{Q} L(w_q)(X_{21}^q + X_{22}^q) \leq m_1 n_2 \sum_{q=1}^{Q} L'(w_q)(X_{21}^q + X_{22}^q) = m_1 n_2 \sum_{q=1}^{Q} L'(w_q)(X_{11}^q + X_{12}^q).$$

Since $L'(w_q) > 0$ by definition and $X_{11}^q + X_{12}^q > 0$ by (14) for every $q = 1, \dots, Q$, it follows that $m_2 n_1 \leq m_1 n_2$, or $\frac{m_2}{m_1} \leq \frac{n_2}{n_1}$.

Note that we can apply the same argument with the roles of G and G' interchanged, and thus get $\frac{n_2}{n_1} \leq \frac{m_2}{m_1}$. This means that $\frac{m_2}{m_1} = \frac{n_2}{n_1}$, so $\frac{m_1}{n_1} = \frac{m_2}{n_2}$, as desired.

Case 2. Now suppose that $\Psi(H)$ has only p_1/c', p_2/c', c/p_1' and c/p_2' vertices.

Let v be a c/p_j' vertex of $\Psi(H)$, where $j = 1$ or $j = 2$. Then all vertices adjacent to v have types p_1/c' or p_2/c'. Let $f_1, \dots f_S$ be all edges incident to v, and let f_s connect v with v_s for $s = 1, \dots, S$ (possibly some of the vertices v_s coincide). Then, by Lemma 9, in the notations of which we have $t = 2$, $t' = n_j + 1$, $r = 2$, $r' = 1$, we obtain

$$\sum_{s=1}^{S} L(v_s) l_v'(f_s) = 2n_j \sum_{s=1}^{S} L'(v_s) l_v'(f_s). \tag{21}$$

Now suppose that $w_1, \dots, w_Q$ are all p_1/c' and p_2/c' vertices of $\Psi(H)$. For every vertex w_q, $q = 1, \dots, Q$, consider all the edges incident to w_q which finish in c/p_1' vertices (note that such edges should always exist), and let $Z_1^q > 0$ be the sum of the l'-labels of these edges at the c/p_1' ends. Analogously, consider all the edges incident to w_q which finish in c/p_2' vertices (note that such edges should always exist), and let $Z_2^q > 0$ be the sum of the l'-labels of these edges at the c/p_2' ends. Let $w_q' = \varphi_*(w_q)$ for $q = 1, \dots, Q$. Then it follows from Lemma 6 that for every $q = 1, \dots, Q$ we have

$$|F_{H', w_q'} : P_{H', w_q'}| = Z_1^q = Z_2^q. \tag{22}$$

Now sum up the equalities of the form (21) for all c/p_1' vertices v, and group the summands with the vertex label corresponding to the same p_i/c' vertex together. We obtain

$$\sum_{q=1}^{Q} L(w_q) Z_1^q = 2n_1 \sum_{q=1}^{Q} L'(w_q) Z_1^q. \tag{23}$$

Analogously, summing the equalities of the form (21) for all c/p_2' vertices v we get

$$\sum_{q=1}^{Q} L(w_q) Z_2^q = 2n_2 \sum_{q=1}^{Q} L'(w_q) Z_2^q. \tag{24}$$

Together with (22), equalities (23) and (24) imply that $n_1 = n_2$.

Note that we can apply the same argument with the roles of G and G' interchanged, and thus get $m_1 = m_2$. It follows that, in particular, $\frac{m_1}{n_1} = \frac{m_2}{n_2}$.

This proves Theorem 3. □

We now prove the main non-commensurability result of this paper.

Theorem 4. *Let $T = T((m_1, 1), \dots, (m_k, 1); 0)$ and $T' = T((n_1, 1), \dots, (n_l, 1); 0)$ for some $k, l \geq 2$. Suppose $G = \mathbb{G}(T)$ and $G' = \mathbb{G}(T')$ are commensurable. Then $M(T)$ and $M(T')$ are commensurable, i.e., $k = l$, and*

$$\frac{n_1}{m_1} = \frac{n_2}{m_2} = \ldots = \frac{n_k}{m_k}. \tag{25}$$

Moreover, if $S = T((m, 2); 0)$ for some $m \geq 2$, then $\mathbb{G}(S)$ and $\mathbb{G}(T')$ are not commensurable.

Proof of Theorem 4. We first prove the first claim. Let c be the central vertex of T and c' be the central vertex of T'. Let $p_1 = p_{1,1}, \dots, p_k = p_{k,1}$ be the pivots of T, and $p'_1 = p'_{1,1}, \dots, p'_l = p'_{l,1}$ be the pivots of T'.

Let $H \leq G$ and $H' \leq G'$ be finite index subgroups such that $H \cong H'$. From Lemma 9 and Lemma 6 applied to both $\Delta = T$ and $\Delta' = T'$ we have a system of equations on the labels of vertices and edges of $\Psi(H)$. We will show that this system of equations does not have any solutions in the positive integers unless $M(T)$ and $M(T')$ are commensurable.

Below we always mean that i ranges between 1 and k, and j ranges between 1 and l, unless stated otherwise.

Note that there could exist the following types of vertices in $\Psi(H)$: c/c', p_i/p'_j, p_i/c', c/p'_j. It is easy to see that the following statements hold:

- Every p_i/p'_j vertex can be connected only with c/c' vertices;
- Every c/c' vertex can be connected only with p_i/p'_j vertices;
- Every p_i/c' vertex can be connected only with c/p'_j vertices;
- Every c/p'_j vertex can be connected only with p_i/c' vertices.

Thus, there are two cases: either $\Psi(H)$ has only p_i/p'_j and c/c' vertices, or $\Psi(H)$ has only p_i/c' and c/p'_j vertices.

Case 1. Suppose first $\Psi(H)$ has only p_i/p'_j and c/c' vertices. We will denote p_i/p'_j vertices as i/j vertices for short.

Let v be an i/j vertex of $\Psi(H)$. Then all vertices adjacent to v have type c/c'. Let $f_1, \dots f_N$ be all edges incident to v, and let f_s connect v with v_s for $s = 1, \dots, S$ (possibly some of the vertices v_s coincide). Then, by Lemma 9, in the notations of which we have $t = m_i + 1, t' = n_j + 1, r = 1, r' = 1$, we obtain

$$m_i \sum_{s=1}^{S} L(v_s) l_v(f_s) = n_j \sum_{s=1}^{S} L'(v_s) l_v(f_s), \tag{26}$$

$$m_i \sum_{s=1}^{S} L(v_s) l'_v(f_s) = n_j \sum_{s=1}^{S} L'(v_s) l'_v(f_s). \tag{27}$$

Now suppose that $w_1, \dots, w_Q$ are all c/c' vertices of $\Psi(H)$. For every vertex w_q, $q = 1, \dots, Q$, consider all the edges incident to w_q which finish in i/j vertices (for fixed i and j), and let X_{ij}^q be the sum of the l-labels of these edges at the i/j ends, and let Y_{ij}^q be the sum of the l'-labels of these edges at the i/j ends; we let $X_{ij}^q = Y_{ij}^q = 0$ if there are no such edges. It follows from Lemma 6 that for all $q = 1, \dots, Q$ we have

$$|F_{H,w_q} : P_{H,w_q}| = X_{i1}^q + X_{i2}^q + \ldots + X_{il}^q$$

for every i, so

$$X_{11}^q + X_{12}^q + \ldots + X_{1l}^q = X_{21}^q + X_{22}^q + \ldots + X_{2l}^q = \ldots = X_{k1}^q + X_{k2}^q + \ldots + X_{kl}^q. \tag{28}$$

Analogously, for all $q = 1, \dots, Q$ if we denote $w'_q = \pi_*(w_q)$, then

$$|F_{H',w'_q} : P_{H',w'_q}| = Y_{1j}^q + Y_{2j}^q + \ldots + Y_{kj}^q$$

for every j, so

$$Y_{11}^q + Y_{21}^q + \ldots + Y_{k1}^q = Y_{12}^q + Y_{22}^q + \ldots + Y_{k2}^q = \ldots = Y_{1l}^q + Y_{2l}^q + \ldots + Y_{kl}^q. \tag{29}$$

Now sum up the equalities of the form (26) for all i/j vertices v (for fixed i and j) and group the summands with the vertex label corresponding to the same c/c' vertex together. We obtain

$$m_i \sum_{q=1}^{Q} L(w_q) X_{ij}^q = n_j \sum_{q=1}^{Q} L'(w_q) X_{ij}^q$$

for all possible i and j. Denote

$$x_{ij} = \sum_{q=1}^{Q} L(w_q) X_{ij}^q, \quad x'_{ij} = \sum_{q=1}^{Q} L'(w_q) X_{ij}^q. \tag{30}$$

Then we get

$$m_i x_{ij} = n_j x'_{ij} \tag{31}$$

for all possible i and j.

Analogously, sum up the equalities of the form (27) for all i/j vertices v (for fixed i and j) and group the summands with the vertex label corresponding to the same c/c' vertex together. We obtain

$$m_i \sum_{q=1}^{Q} L(w_q) Y_{ij}^q = n_j \sum_{q=1}^{Q} L'(w_q) Y_{ij}^q$$

for all possible i and j. Denote

$$y_{ij} = \sum_{q=1}^{Q} L(w_q) Y_{ij}^q, \quad y'_{ij} = \sum_{q=1}^{Q} L'(w_q) Y_{ij}^q. \tag{32}$$

Then we get

$$m_i y_{ij} = n_j y'_{ij} \tag{33}$$

for all possible i and j.

From (28) and (30) we get

$$x_{11} + x_{12} + \ldots + x_{1l} = x_{21} + x_{22} + \ldots + x_{2l} = \ldots = x_{k1} + x_{k2} + \ldots + x_{kl}, \tag{34}$$

$$x'_{11} + x'_{12} + \ldots + x'_{1l} = x'_{21} + x'_{22} + \ldots + x'_{2l} = \ldots = x'_{k1} + x'_{k2} + \ldots + x'_{kl}, \tag{35}$$

for all possible i and j. Summing the equalities (31) with the same i we get the following system of equations

$$\begin{cases} m_1(x_{11} + x_{12} + \ldots + x_{1l}) = n_1x'_{11} + n_2x'_{12} + \ldots + n_lx'_{1l} \\ m_2(x_{21} + x_{22} + \ldots + x_{2l}) = n_1x'_{21} + n_2x'_{22} + \ldots + n_lx'_{2l} \\ \quad \ldots \\ m_{k-1}(x_{k-1,1} + x_{k-1,2} + \ldots + x_{k-1,l}) = n_1x'_{k-1,1} + n_2x'_{k-1,2} + \ldots + n_lx'_{k-1,l} \\ m_k(x_{k1} + x_{k2} + \ldots + x_{kl}) = n_1x'_{k1} + n_2x'_{k2} + \ldots + n_lx'_{kl} \end{cases} \tag{36}$$

Analogously, from (29) and (32) we get

$$y_{11} + y_{21} + \ldots + y_{k1} = y_{12} + y_{22} + \ldots + y_{k2} = \ldots = y_{1l} + y_{2l} + \ldots + y_{kl}, \tag{37}$$

$$y'_{11} + y'_{21} + \ldots + y'_{k1} = y'_{12} + y'_{22} + \ldots + y'_{k2} = \ldots = y'_{1l} + y'_{2l} + \ldots + y'_{kl}, \tag{38}$$

for all possible i and j. Summing the equalities (33) with the same j we get the following system of equations

$$\begin{cases} m_1y_{11} + m_2y_{21} + \ldots + m_ky_{k1} = n_1(y'_{11} + y'_{21} + \ldots + y'_{k1}) \\ m_1y_{12} + m_2y_{22} + \ldots + m_ky_{k2} = n_2(y'_{12} + y'_{22} + \ldots + y'_{k2}) \\ \quad \ldots \\ m_1y_{1,l-1} + m_2y_{2,l-1} + \ldots + m_ky_{k,l-1} = n_1(y'_{1,l-1} + y'_{2,l-1} + \ldots + y'_{k,l-1}) \\ m_1y_{1l} + m_2y_{2l} + \ldots + m_ky_{kl} = n_1(y'_{1l} + y'_{2l} + \ldots + y'_{kl}) \end{cases} \tag{39}$$

Let $A = (a_{ij})$ be the matrix with k rows and l columns, such that $a_{ij} = 1$ if there are i/j vertices in $\Psi(H)$ and $a_{ij} = 0$ otherwise. Note that for every $q = 1, \ldots, Q$ we have $L(w_q) > 0$, $L'(w_q) > 0$, and $X^q_{ij} = 0$ if and only $Y^q_{ij} = 0$, if and only if there are no vertices of type i/j adjacent to w_q. Obviously every vertex of type i/j of $\Psi(H)$ should be adjacent to at least one c/c' vertex, i.e. to at least one of $w_1, \ldots, w_Q$. This means that $x_{ij} \geq 0$, $x'_{ij} \geq 0$, and $x_{ij} = 0$ if and only if $x'_{ij} = 0$, if and only if $X^q_{ij} = 0$ for all $q = 1, \ldots, Q$, if and only if there are no vertices of type i/j in $\Psi(H)$, if and only if $a_{ij} = 0$. Analogously $y_{ij} \geq 0$, $y'_{ij} \geq 0$, and $y_{ij} = 0$ if and only if $y'_{ij} = 0$, if and only if $Y^q_{ij} = 0$ for all $q = 1, \ldots, Q$, if and only if there are no vertices of type i/j in $\Psi(H)$, if and only if $a_{ij} = 0$.

Note also that for every i there should exist some j such that there are i/j vertices in $\Psi(H)$. Analogously, for every j there should exist some i such that i/j vertices exist in $\Psi(H)$. This means that the matrix A does not have zero rows or columns.

Combining (34), (35), (36), (37), (38), (39), we have the following equations and conditions on $x_{ij}, x'_{ij}, y_{ij}, y'_{ij}, a_{ij}$:

$$\begin{cases} m_1(x_{11}+x_{12}+\ldots+x_{1l}) = n_1x'_{11}+n_2x'_{12}+\ldots+n_lx'_{1l} \\ m_2(x_{21}+x_{22}+\ldots+x_{2l}) = n_1x'_{21}+n_2x'_{22}+\ldots+n_lx'_{2l} \\ \qquad \ldots \\ m_{k-1}(x_{k-1,1}+x_{k-1,2}+\ldots+x_{k-1,l}) = n_1x'_{k-1,1}+n_2x'_{k-1,2}+\ldots+n_lx'_{k-1,l} \\ m_k(x_{k1}+x_{k2}+\ldots+x_{kl}) = n_1x'_{k1}+n_2x'_{k2}+\ldots+n_lx'_{kl} \\ x_{11}+x_{12}+\ldots+x_{1l} = x_{21}+x_{22}+\ldots+x_{2l} = \ldots = x_{k1}+x_{k2}+\ldots+x_{kl} \\ x'_{11}+x'_{12}+\ldots+x'_{1l} = x'_{21}+x'_{22}+\ldots+x'_{2l} = \ldots = x'_{k1}+x'_{k2}+\ldots+x'_{kl} \end{cases} \tag{40}$$

$$\begin{cases} m_1y_{11}+m_2y_{21}+\ldots+m_ky_{k1} = n_1(y'_{11}+y'_{21}+\ldots+y'_{k1}) \\ m_1y_{12}+m_2y_{22}+\ldots+m_ky_{k2} = n_2(y'_{12}+y'_{22}+\ldots+y'_{k2}) \\ \qquad \ldots \\ m_1y_{1,l-1}+m_2y_{2,l-1}+\ldots+m_ky_{k,l-1} = n_{l-1}(y'_{1,l-1}+y'_{2,l-1}+\ldots+y'_{k,l-1}) \\ m_1y_{1l}+m_2y_{2l}+\ldots+m_ky_{kl} = n_l(y'_{1l}+y'_{2l}+\ldots+y'_{kl}) \\ y_{11}+y_{21}+\ldots+y_{k1} = y_{12}+y_{22}+\ldots+y_{k2} = \ldots = y_{1l}+y_{2l}+\ldots+y_{kl} \\ y'_{11}+y'_{21}+\ldots+y'_{k1} = y'_{12}+y'_{22}+\ldots+y'_{k2} = \ldots = y'_{1l}+y'_{2l}+\ldots+y'_{kl} \end{cases} \tag{41}$$

$$x_{ij}=0 \Leftrightarrow x'_{ij}=0 \Leftrightarrow y_{ij}=0 \Leftrightarrow y'_{ij}=0 \Leftrightarrow a_{ij}=0, \quad i=1,\ldots k,\ j=1,\ldots l \tag{42}$$

$$\forall i\,\forall j\ x_{ij}, x'_{ij}, y_{ij}, y'_{ij} \geq 0, \quad \forall i\,\exists j : x_{ij}>0, \quad \forall j\,\exists i : x_{ij}>0, \quad i=1,\ldots k,\ j=1,\ldots l. \tag{43}$$

Without loss of generality, we can assume that $k \geq l$. By $[\alpha]$ we denote the (lower) integer part of α, so $[l/2] = l/2$ if l is even, and $[l/2] = (l-1)/2$ if l is odd.

We first claim that $a_{ij} = a_{k+1-i,l+1-j} = 0$ if $i \leq [l/2]$ or $j \leq [l/2]$, provided $i \neq j$. We prove this claim by induction on d_{ij}, where $d_{ij} = min\,(i,\ j)$. Thus we are interested in the case when $1 \leq d_{ij} \leq [l/2]$.

The base of induction is the case when $d_{ij} = 1$, so either $i = 1$, or $j = 1$. This case is included in the inductive step below, with $r = 1$.

Suppose the claim is proved for $d(i,j) \leq r-1$, where $1 \leq r \leq [l/2]$, and we want to prove it for $d(i,j) = r$. We have $a_{ij} = 0$ if $i \leq r-1$ or $j \leq r-1$, provided $i \neq j$, and $a_{ij} = 0$ if $i \geq k-r+2$ or $j \geq l-r+2$, provided $k-i \neq l-j$ (in the case $r=1$ there are no conditions). By (42), this means that the r-th equation of (40) has the following form

$$m_r(x_{rr}+x_{r,r+1}+\ldots+x_{r,l+1-r}) = n_rx'_{rr}+n_{r+1}x'_{r,r+1}+\ldots+n_{l+1-r}x'_{r,l+1-r}, \tag{44}$$

and the $(k+1-r)$-th equation of (40) has the following form

$$\begin{aligned} m_{k+1-r}(x_{k+1-r,r}+x_{k+1-r,r+1}+\ldots+x_{k+1-r,l+1-r}) = \\ = n_rx'_{k+1-r,r}+n_{r+1}x'_{k+1-r,r+1}+\ldots+n_{l+1-r}x'_{k+1-r,l+1-r}. \end{aligned} \tag{45}$$

Also from the last two lines of equations of (40) we obtain

$$x_{rr}+x_{r,r+1}+\ldots+x_{r,l+1-r} = x_{k+1-r,r}+x_{k+1-r,r+1}+\ldots+x_{k+1-r,l+1-r}, \tag{46}$$

$$x'_{rr} + x'_{r,r+1} + \ldots + x'_{r,l+1-r} = x'_{k+1-r,r} + x'_{k+1-r,r+1} + \ldots + x'_{k+1-r,l+1-r}. \tag{47}$$

Since $n_r < n_{r+1} < \ldots < n_{l+1-r}$, (44) implies

$$m_r(x_{rr} + x_{r,r+1} + \ldots + x_{r,l+1-r}) \geq n_r(x'_{rr} + x'_{r,r+1} + \ldots + x'_{r,l+1-r}) \tag{48}$$

and equality in (48) is obtained if and only if $x'_{r,r+1} = x'_{r,r+2} = \ldots = x'_{r,l+1-r} = 0$.

Also (45) implies

$$\begin{aligned} m_{k+1-r}(x_{k+1-r,r} + x_{k+1-r,r+1} + \ldots + x_{k+1-r,l+1-r}) \leq \\ \leq n_{l+1-r}(x'_{k+1-r,r} + x'_{k+1-r,r+1} + \ldots + x'_{k+1-r,l+1-r}), \end{aligned} \tag{49}$$

and equality in (49) is obtained if and only if $x'_{k+1-r,r} = x'_{k+1-r,r+1} = \ldots = x'_{k+1-r,l-r} = 0$. Multiplying (48) by m_{k+1-r}, (49) by m_r and using (46), (47), we obtain

$$\begin{aligned} m_{k+1-r}n_r(x'_{rr} + x'_{r,r+1} + \ldots + x'_{r,l+1-r}) \leq m_{k+1-r}m_r(x_{rr} + x_{r,r+1} + \ldots + x_{r,l+1-r}) = \\ = m_r m_{k+1-r}(x_{k+1-r,r} + x_{k+1-r,r+1} + \ldots + x_{k+1-r,l+1-r}) \leq \\ \leq m_r n_{l+1-r}(x'_{k+1-r,r} + x'_{k+1-r,r+1} + \ldots + x'_{k+1-r,l+1-r}) = \\ = m_r n_{l+1-r}(x'_{rr} + x'_{r,r+1} + \ldots + x'_{r,l+1-r}). \end{aligned} \tag{50}$$

Note that by (42), (43) we have $x'_{rr} + x'_{r,r+1} + \ldots + x'_{r,l+1-r} \neq 0$ (since otherwise the r-th row of A is zero, which is impossible). So (50) implies

$$m_{k+1-r}n_r \leq m_r n_{l+1-r}. \tag{51}$$

Now apply analogous arguments to (41). By induction hypothesis the r-th equation of (41) has the following form

$$m_r y_{rr} + m_{r+1}y_{r+1,r} + \ldots + m_{k+1-r}y_{k+1-r,r} = n_r(y'_{rr} + y'_{r+1,r} + \ldots + y'_{k+r-1,r}), \tag{52}$$

and the $(l+1-r)$-th equation of (41) has the following form

$$\begin{aligned} m_r y_{r,l+1-r} + m_{r+1}y_{r+1,l+1-r} + \ldots + m_{k+1-r}y_{k+1-r,l+1-r} = \\ = n_{l+1-r}(y'_{r,l+1-r} + y'_{r+1,l+1-r} + \ldots + y'_{k+r-1,l+1-r}). \end{aligned} \tag{53}$$

Also from the last two lines of equations (41) we obtain

$$y_{rr} + y_{r+1,r} + \ldots + y_{k+1-r,r} = y_{r,l+1-r} + y_{r+1,l+1-r} + \ldots + y_{k+1-r,l+1-r}, \tag{54}$$

$$y'_{rr} + y'_{r+1,r} + \ldots + y'_{k+1-r,r} = y'_{r,l+1-r} + y'_{r+1,l+1-r} + \ldots + y'_{k+1-r,l+1-r}. \tag{55}$$

Since $m_r < m_{r+1} < \ldots < m_{k+1-r}$, (52) implies

$$m_r(y_{rr} + y_{r+1,r} + \ldots + y_{k+1-r,r}) \leq n_r(y'_{rr} + y'_{r+1,r} + \ldots + y'_{k+r-1,r}), \tag{56}$$

and equality in (56) is obtained if and only if $y_{r+1,r} = y_{r+2,r} = \ldots = y_{k+1-r,r} = 0$.

Also (53) implies

$$\begin{aligned} m_{k+1-r}(y_{r,l+1-r} + y_{r+1,l+1-r} + \ldots + y_{k+1-r,l+1-r}) \geq \\ \geq n_{l+1-r}(y'_{r,l+1-r} + y'_{r+1,l+1-r} + \ldots + y'_{k+r-1,l+1-r}), \end{aligned} \tag{57}$$

and equality in (57) is attained if and only if $y_{r,l+1-r} = y_{r+1,l+1-r} = \ldots = y_{k-r,l+1-r} = 0$. Multiplying (56) by n_{l+1-r}, (57) by n_r and using (54), (55), we obtain

$$\begin{aligned}(58)\quad n_{l+1-r}m_r(y_{rr} + y_{r+1,r} + \ldots + y_{k+1-r,r}) &\leq n_{l+1-r}n_r(y'_{rr} + y'_{r+1,r} + \ldots + y'_{k+r-1,r}) = \\ &= n_r n_{l+1-r}(y'_{r,l+1-r} + y'_{r+1,l+1-r} + \ldots + y'_{k+r-1,l+1-r}) \leq \\ &\leq n_r m_{k+1-r}(y_{r,l+1-r} + y_{r+1,l+1-r} + \ldots + y_{k+1-r,l+1-r}) = \\ &= n_r m_{k+1-r}(y_{rr} + y_{r+1,r} + \ldots + y_{k+1-r,r}).\end{aligned}$$

Note that by (42), (43) we have $y_{rr} + y_{r+1,r} + \ldots + y_{k+1-r,r} \neq 0$ (since otherwise the r-th column of A is zero, which is impossible). So (58) implies

$$n_{l+1-r}m_r \leq n_r m_{k+1-r}.$$

Combined with (51), this gives

$$(59)\qquad m_{k+1-r}n_r = m_r n_{l+1-r},$$

so all the inequalities in (50), (58), and then also in (48), (49), (56), (57), turn into equalities. As mentioned above, this means that

$$x'_{r,r+1} = x'_{r,r+2} = \ldots = x'_{r,l+1-r} = 0, \quad x'_{k+1-r,r} = x'_{k+1-r,r+1} = \ldots = x'_{k+1-r,l-r} = 0,$$

so by (42) we have

$$a_{r,r+1} = a_{r,r+2} = \ldots = a_{r,l+1-r} = 0, \quad a_{k+1-r,r} = a_{k+1-r,r+1} = \ldots = a_{k+1-r,l-r} = 0,$$

and

$$y_{r+1,r} = y_{r+2,r} = \ldots = y_{k+1-r,r} = 0, \quad y_{r,l+1-r} = y_{r+1,l+1-r} = \ldots = y_{k-r,l+1-r} = 0,$$

so by (42) we have

$$a_{r+1,r} = a_{r+2,r} = \ldots = a_{k+1-r,r} = 0, \quad a_{r,l+1-r} = a_{r+1,l+1-r} = \ldots = a_{k-r,l+1-r} = 0.$$

Combined with the induction hypothesis, this proves that $a_{ij} = a_{k+1-i,l+1-j} = 0$ if $i \leq r$ or $j \leq r$, provided $i \neq j$. This proves the claim.

Thus we have proved that

$$(60)\qquad a_{ij} = a_{k+1-i,l+1-j} = 0, \text{ if } i \leq [l/2] \text{ or } j \leq [l/2], \text{ provided } i \neq j.$$

We now show that $k = l$. Suppose, on the contrary, that $k > l$. Suppose first l is even. Then we have from (60) that $a_{ij} = 0$ if $j \leq l/2$ and $i \neq j$, and $a_{ij} = 0$ if $j \geq l/2 + 1$ and $k - i \neq l - j$. In particular, taking $i = l/2 + 1$, we get that $a_{l/2+1,j} = 0$ for $j \leq l/2$ (since for these pairs $i \neq j$), and $a_{l/2+1,j} = 0$ for $j \geq l/2 + 1$ (since for these pairs $k - i = k - l/2 - 1 > l/2 - 1 \geq l - j$), so A has a zero row, a contradiction.

So l is odd. Then it follows from (60) that $a_{ij} = 0$ if $j \leq (l-1)/2$ and $i \neq j$, and $a_{ij} = 0$ if $j \geq (l+3)/2$ and $k - i \neq l - j$. In particular, $a_{(l+1)/2,j} = a_{(l+3)/2,j} = 0$ when $j \neq (l+1)/2$. Denote $(l+1)/2 = b$, then $(l+3)/2 = b+1$. Due to (42), this means that among the equations (40) we have the following:

$$\begin{cases} m_b x_{bb} = n_b x'_{bb}, \\ m_{b+1} x_{b+1,b} = n_b x'_{b+1,b}, \\ x_{bb} = x_{b+1,b}, \\ x'_{bb} = x'_{b+1,b}. \end{cases}$$

This immediately implies $m_b = m_{b+1}$, a contradiction.

This shows that $k = l$. Then it is easy to see that it follows from (60) that $a_{ij} = 0$ if $i \neq j$. Then A is the identity matrix, since it does not have zero rows or columns, so, according to (42), (43), we have $x_{ij} = x'_{ij} = 0$ if $i \neq j$, and $x_{ii} > 0$ for all $i = 1, \dots k$, therefore (40) turns into

$$\begin{cases} m_1 x_{11} = n_1 x'_{11} \\ m_2 x_{22} = n_2 x'_{22} \\ \quad \dots \\ m_k x_{kk} = n_k x'_{kk} \\ x_{11} = x_{22} = \dots = x_{kk} \\ x'_{11} = x'_{22} = \dots = x'_{kk}, \end{cases}$$

which immediately implies (25). This finishes Case 1.

Case 2. Now suppose that $\Psi(H)$ has only p_i/c' and c/p'_j vertices, $i = 1, \dots, k$, $j = 1, \dots, l$. The proof in this case is almost the same as the proof of Case 2 in Theorem 3.

Let v be a c/p'_j vertex of $\Psi(H)$, $j = 1, \dots, l$. Then all vertices adjacent to v have types p_i/c', $i = 1, \dots, k$. Let $f_1, \dots f_S$ be all edges incident to v, and let f_s connect v with v_s for $s = 1, \dots, S$ (possibly some of the vertices v_s coincide). Then, by Lemma 9, in the notations of which we have $t = k$, $t' = n_j + 1$, $r = k$, $r' = 1$, we obtain

$$\sum_{s=1}^{S} L(v_s) l'_v(f_s) = \frac{k}{k-1} n_j \sum_{s=1}^{S} L'(v_s) l'_v(f_s). \tag{61}$$

Now suppose that $w_1, \dots, w_Q$ are all p_i/c' vertices of $\Psi(H)$, for all $i = 1, \dots, k$. For every vertex w_q, $q = 1, \dots, Q$ and $j = 1, 2$, consider all the edges incident to w_q which finish in c/p'_j vertices (note that such edges should always exist), and let Z_j^q be the sum of the l'-labels of these edges at the c/p'_j ends. Then we have $Z_j^q > 0$ for every $q = 1, \dots, Q$ and $j = 1, 2$. Let $w'_q = \varphi_*(w_q)$ for $q = 1, \dots, Q$. Then it follows from Lemma 6 that for every $q = 1, \dots, Q$ we have

$$|F_{H', w'_q} : P_{H', w'_q}| = Z_1^q = Z_2^q. \tag{62}$$

Now sum up the equalities of the form (61) for all c/p'_1 vertices v, and group the summands with the vertex label corresponding to the same p_i/c' vertex together. We obtain

$$\sum_{q=1}^{Q} L(w_q) Z_1^q = \frac{k}{k-1} n_1 \sum_{q=1}^{Q} L'(w_q) Z_1^q. \tag{63}$$

Analogously, summing the equalities of the form (61) for all c/p'_2 vertices v we get

$$\sum_{q=1}^{Q} L(w_q) Z_2^q = \frac{k}{k-1} n_2 \sum_{q=1}^{Q} L'(w_q) Z_2^q. \tag{64}$$

Together with (62), equalities (63) and (64) imply that $n_1 = n_2$, a contradiction.

This finishes the proof of the first claim of the theorem.

The proof of the second claim is similar, but easier. Indeed, let $m_1 = m_2 = m$, $k = 2$, and apply the arguments as in the proof of the first claim above. In Case 1, it is easy to see by the argument as above that (59) still holds with $r = 1$, so $m_2 n_1 = m_1 n_l$, but $m_1 = m_2$, so $n_1 = n_l$, a contradiction. The proof of Case 2 is the same as above.

This proves Theorem 4. □

3. Characterisation of commensurability classes of RAAGs defined by trees of diameter 4

We now turn to prove commensurability of some RAAGs defined by trees of diameter 4.

Proposition 1. *Let $\mathbb{G} = \mathbb{G}(T)$, where $T = T((v_1, 1), \dots, (v_l, 1); 0)$, $l \geq 2$. For any $k_1, \dots, k_l$ there exist a non-negative integer q and a finite index subgroup H of $\mathbb{G}$ such that $H \simeq \mathbb{G}(S)$, where S is a tree of diameter 4 of the form $S = T((v_1, k_1), \dots, (v_l, k_l); q)$. In particular, $\mathbb{G}(S)$ and $\mathbb{G}(T)$ are commensurable.*

If $\mathbb{G} = \mathbb{G}(T)$, where $T = T((d, 2); 0)$, then for any $k > 2$ there exists a finite index subgroup H of $\mathbb{G}$ such that $H \simeq \mathbb{G}(S)$, where $S = T((d, k); 0)$.

Proof. We first consider the case when $l = 1$, i.e. $T = T((d, 2); 0)$. Let p_1, p_2 be the pivots of T. Let ϕ be the epimorphism $\mathbb{G} \to \mathbb{Z}_k$, induced by the map $p_1 \mapsto 1$ and $x \mapsto 0$, where x is any canonical generator of $\mathbb{G}$ different from p_1. It is not hard to see that $\ker \phi$ is isomorphic to $\mathbb{G}(S)$, see [KK13, BN08].

Suppose now $l \geq 2$. Denote the only pivot of valency v_i in T by p_i, $i = 1, \dots, l$, and the central vertex of T by c. Let $V(T) = X$. Let $F(X)$ be the free group with the basis X. Then there is a natural epimorphism $\phi : F(X) \to \mathbb{G}$. Fix $k_1, \dots, k_l$ and without loss of generality assume that $k_1 \leq k_2 \leq \dots \leq k_l$. We can always suppose that $k_l > 1$, since otherwise $k_1 = k_2 = \dots = k_l = 1$, and the claim is obvious. Let $x_{i,1}, \dots, x_{i,v_i}$ be all leaves of T adjacent to the pivot p_i, $i = 1, \dots, l$.

Consider the finite index subgroup $G < F(X)$ defined as the subgroup corresponding to the finite cover of the bouquet of $|X|$ circles K defined as follows. Below by $(a_1, \dots, a_{k_l})$ we mean a simple cycle of length k_l, with vertices appearing in the order $a_1, \dots, a_{k_l}, a_1$, and analogously with other cycles. We take two cycles: the first one of the form $(a_1, \dots, a_{k_l})$, of length k_l, with all edges labelled by p_1; and the second one of the form $(b_1, \dots, b_{k_1})$, of length k_1, with all edges labelled by p_l; we identify these cycles by a vertex, $a_1 = b_1$. This vertex is the basepoint of the based cover we construct. The degree of the finite cover of the bouquet of $|X|$ circles we construct is $k_1 + k_l - 1$, so no new vertices will be added, only edges. We first complete the constructed graph to a cover of the free group $F(p_1, p_l)$, by adding loops labelled by p_1 at vertices $b_2, \dots, b_{k_1}$ (no loops are added if $k_1 = 1$), and adding loops labelled by p_l at vertices $a_2, \dots, a_{k_l}$.

If $l = 2$, then we are done with the construction of the cover. If $l > 2$, we consider two cases. In the first case we suppose that $k_2 = 1$, so for some $2 \leq j < l$ we have $1 = k_1 = k_2 = \dots = k_j < k_{j+1} \leq k_{j+2} \leq \dots \leq k_l$. In the second case we assume that $k_2 > 1$.

In the first case, for every $i = 2, \dots, j$, we add a cycle $(a_1, \dots, a_{k_l})$ labelled by p_i. For every $i = j + 1, \dots, l-1$, we add loops labelled by p_i to vertices $a_1, \dots, a_{k_i - 1}$, and add a cycle $(a_{k_i}, a_{k_i+1}, \dots, a_{k_l})$, of length $k_l - k_i + 1$, with all edges labelled by p_i, see Figure 3. Adding loops labelled by the



FIGURE 3. Constructing the cover, case 1

generators from $X \setminus \{p_1, \dots, p_l\}$ at all vertices of the graph we obtain a finite cover of the bouquet of $|X|$ circles and hence this defines a finite index subgroup G of $F(X)$.



FIGURE 4. Constructing the cover, case 2

In the second case, for every $i = 2, \dots, l-1$, we do the following. First add a cycle $(b_1, \dots, b_{k_1})$ with all edges labelled by p_i. Then add loops labelled by p_i to vertices $a_2, \dots, a_{k_i - 1}$. Finally, add a cycle $(a_{k_i}, a_{k_i+1}, \dots, a_{k_l})$, of length $k_l - k_i + 1$, labelled by p_i, see Figure 4. Adding loops labelled by the generators $X \setminus \{p_1, \dots, p_l\}$ at all vertices of the graph we obtain a finite cover of the bouquet of $|X|$ circles and hence this defines a finite index subgroup G of $F(X)$.

We continue under the assumptions of the second case. Proof in the first case is analogous.

Choosing the maximal subtree spanned by the edges $\{(a_i, a_{i+1}), i = 1, \dots, k_l - 1\}$, labelled by p_l, and $\{(b_j, b_{j+1}) \mid \ j = 1, \dots, k_1 - 1\}$, labelled by p_1, and using Tietze transformations, it is not difficult to see that $\{p_i^{k_1}, \left(p_i^{k_l - k_i + 1}\right)^{p_1^{k_i - 1}} \mid i = 2, \dots, l-1\}$ is part of the generating set of G, which consists of the following generators:

A: • $\{p_i^{k_1}, p_i^{(p_1^{j_i - 1})}, \left(p_i^{k_l - k_i + 1}\right)^{p_1^{k_i - 1}} \mid j_i = 2, \dots, k_i - 1,\ i = 2, \dots, l-1\}$;

- $\{p_1^{k_l}, p_1^{(p_l^{j_1-1})} \mid j_1 = 2, \dots, k_1\}$;
- $\{p_l^{k_1}, p_l^{(p_1^{j_l-1})} \mid j_l = 2, \dots, k_l\}$.

B: $\{x_{i,m_i}^{p_1^r} \mid r = 0, \dots, k_l - 1\}$, $i = 1, \dots, l$, $m_i = 1, \dots, v_i$;

$\{x_{i,m_i}^{p_l^s} \mid s = 1, \dots, k_1 - 1\}$, $i = 1, \dots, l$, $m_i = 1, \dots, v_i$.

C: $\{c^{p_1^r} \mid r = 0, \dots, k_l - 1\}$, $\{c^{p_l^s} \mid s = 1, \dots, k_1 - 1\}$.

We have epimorphisms $\phi : F(X) \to \mathbb{G}$ and $\psi : \mathbb{G} \to F(p_1, \dots, p_l)$, where $F(p_1, \dots, p_l)$ is the free group with the basis $p_1, \dots, p_l$. By construction, the image of $\psi(\phi(G))$ has index $k_1 + k_l - 1$ in $F(p_1, \dots, p_l)$. Hence,

$$[F(X) : G] \geq [\mathbb{G} : \phi(G)] \geq [F(p_1, \dots, p_l) : \psi(\phi(G))],$$

and it follows that $H = \phi(G)$ has index $k_1 + k_l - 1$ in $\mathbb{G}$. Thus H is a finite index subgroup of $\mathbb{G}$, and H is generated by the set $\phi(A) \cup \phi(B) \cup \phi(C)$. Note that in $\mathbb{G}$ all elements of C become equal to c, so $\phi(C)$ consists just of one generator c.

We now show that $H \simeq \mathbb{G}(S)$, where $S = T((v_1, k_1), \dots, (v_l, k_l); q)$. Note that the group $\mathbb{G}$ splits as a fundamental group of a star of groups: $\mathbb{G} = \pi_1(\mathbb{B})$, with l leaves, with vertex groups at leaves equal to $B_i = \langle c, p_i, x_{i,1}, \dots, x_{i,v_i} \rangle$, $i = 1, \dots, l$, and vertex group at the center vertex equal to $B_c = \langle c, p_1, \dots, p_l \rangle$. The edge groups are $E_i = \langle c, p_i \rangle$, $i = 1, \dots, l$. This is the reduced centralizer splitting of $\mathbb{G}$. Let $\mathbf{T}$ be the Bass-Serre tree corresponding to this splitting of $\mathbb{G}$.

Recall that by Bass-Serre theory vertices of $\mathbf{T}$ correspond to left cosets by the vertex groups of $\mathbb{B}$, and edges of $\mathbf{T}$ correspond to left cosets by the edge groups of $\mathbb{B}$. Consider the finite subgraph Y_0 of the tree $\mathbf{T}$, which consists of the vertex B_c and the incident edges $p_1^{j_i} E_i$ going to the vertices $p_1^{j_i} B_i$, $j_i = 0, \dots, k_i - 1$, $i = 2, \dots, l$, and edges $p_l^{j_1} E_1$ going to the vertices $p_l^{j_1} B_1$, $j_1 = 0, \dots, k_1 - 1$, together with these vertices. Note that all these vertices are indeed different. Thus Y_0 as a graph is a star with $k_1 + \dots + k_l$ leaves. We claim that Y_0 is the fundamental domain under the action of H on $\mathbf{T}$, i.e., Y_0 contains exactly one representative of each vertex and edge orbit under the action of H on $\mathbf{T}$.

First we show that Y_0 cannot contain two vertices or edges in the same orbit under the action of H. It suffices to show this for vertices of Y_0. It is obvious that B_c is not in the same H-orbit with any other vertex of Y_0, since it is not even in the same $\mathbb{G}$-orbit. Suppose that $u = p_1^{j_i} B_i$ is in the same H-orbit with some other vertex v of Y_0, then v can only be of the form $v = p_1^{j_i'} B_i$, where $j_i' \neq j_i$; here $j_i, j_i' = 0, \dots, k_i - 1$, $i = 2, \dots, l$. Then there exists some $d \in B_i = \langle c, p_i, x_{i,1}, \dots, x_{i,v_i} \rangle$ such that

$$H p_1^{j_i} d = H p_1^{j_i'}.$$

If $j_i < k_i - 1$, then from the definition of H, see Figure 4, we have that

$$H p_1^{j_i} B_i = H p_1^{j_i},$$

and it follows that $H p_1^{j_i'} = H p_1^{j_i}$, so $p_1^{j_i' - j_i} \in H$, which is a contradiction by definition of H.

If $j_i = k_i - 1$, then $j_i' < k_i - 1$, and we get a contradiction in the same way as above.

In the same way we can prove that the vertex $p_l^{j_1}B_1$, $j_1 = 0, \dots, k_1 - 1$, is not in the same H-orbit with some other vertex in Y_0. Thus indeed Y_0 does not contain two vertices or edges in the same orbit under the action of H.

Now we show that Y_0 contains at least one representative of each orbit, and so Y_0 is a fundamental domain. Note that it suffices to show that every edge of $\mathbf{T}$ which is incident to some edge in Y_0 can be taken to some edge in Y_0 by an element of H. Let e be some edge of $\mathbf{T}$ incident to the vertex B_c, then by Bass-Serre theory $e = gE_i$, where $g \in B_c$, and we can suppose g does not contain c, so $g = g(p_1, \dots p_l)$. Suppose $i > 1$, the case $i = 1$ is analogous. By definition of H, see Figure 4,

$$Q = \{p_1^s,\ s = 0, \dots, k_i - 1;\ p_1^{k_i-1}p_i^m,\ m = 1, \dots, k_l - k_i;\ p_l^t,\ t = 1, \dots k_1 - 1\}$$

is the set of coset representatives of H in $\mathbb{G}$, so $g = hq$ for some $q \in Q$, and $e = gE_i$ is in the same H-orbit as qE_i. Since $p_i \in E_i$, by definition of Y_0 the edge qE_i is in Y_0, and so we are done.

Now, if f is some edge of $\mathbf{T}$ incident to the vertex $p_1^{j_i}B_i$, $j_i = 0, \dots, k_i - 1$, $i = 2, \dots, l$, then by Bass-Serre theory $f = gE_i$, where $g = p_1^{j_i}b$, $b \in B_i$, so $f = p_1^{j_i}bE_i$, and we can suppose b does not contain p_i, so $b = w(c, x_{i,1}, \dots, x_{i,v_i})$. Then, by definition of H, $p_1^{j_i}b(p_1^{j_i})^{-1} = h \in H$, so $f = p_1^{j_i}bE_i = hp_1^{j_i}E_i$ is in the same H-orbit as $p_1^{j_i}E_i$, which is in Y_0 by definition. The case when f is some edge of $\mathbf{T}$ incident to the vertex $p_l^{j_1}B_1$, $j_1 = 0, \dots, k_1 - 1$ is analogous.

This proves that Y_0 is the fundamental domain under the action of H on $\mathbf{T}$. Let $H \cong \pi_1(\mathbb{Y})$ be the induced splitting of H as a fundamental group of a graph of groups, corresponding to the induced action of H on $\mathbf{T}$. Let Y be the underlying graph of $\mathbb{Y}$. Let $\pi : \mathbf{T} \to Y$ be the natural projection morphism.

It follows that the morphism π restricted to Y_0 induces an isomorphism of graphs $\pi_{Y_0} : Y_0 \to Y$, and the vertex group at $\pi(v)$ is equal to the stabilizer of the vertex v for every vertex v of Y_0. This means that $\mathbb{Y}$ is a star with $k_1 + \dots + k_l$ leaves, $u_{1,1}, \dots, u_{1,k_1}; \dots; u_{l,1}, \dots, u_{l,k_l}$ and the center vertex z. The vertex group at z is $H \cap \langle c, p_1, \dots, p_l\rangle = \langle \text{generators of type } A, c\rangle$. The vertex groups $G_{u_{i,j_i}}$ at leaves u_{i,j_i} are the following:

$$\begin{cases} G_{u_{i,j_i}} = H \cap B_i^{(p_1^{j_i})} = H \cap \langle c, p_i, x_{i,1}, \dots, x_{i,v_i}\rangle^{p_1^{j_i}}, & j_i = 0, \dots, k_i - 1,\ i = 2, \dots, l; \\ G_{u_{1,j_1}} = H \cap B_1^{(p_l^{j_1})} = H \cap \langle c, p_1, x_{1,1}, \dots, x_{1,v_1}\rangle^{p_l^{j_1}}, & j_1 = 0, \dots, k_1 - 1. \end{cases}$$

The edge groups of $\mathbb{Y}$ are the corresponding intersections of vertex groups. Since $c \in H$ it follows that the vertex groups are direct products of the free group and the infinite cyclic group. Moreover, since, by construction, conjugates of $x_{i,1}, \dots, x_{i,v_i-1}$ belong to H, it follows that every edge group of $\mathbb{Y}$ is the direct product of the cyclic group generated by c and a subgroup of $\langle p_1, \dots, p_l\rangle$.

Suppose $S = T((v_1, k_1), \dots, (v_l, k_l); q)$ has pivots $p_{i,1}, \dots, p_{i,k_i}$ of valency $v_i + 1$, $i = 1, \dots, l$. Let $x_{i,j_i,1}, \dots, x_{i,j_i,v_i}$ be all leaves of S which are adjacent to the pivot p_{i,j_i}, for $j_i = 1, \dots, k_i$, $i = 1, \dots, l$. Denote the center of S by c', and the hairs of S by $y_1, \dots, y_q$.

Since Y is a tree, $\pi_1(\mathbb{Y})$ is a free product with amalgamation and it is easy to show that there is an isomorphism $\pi_1(\mathbb{Y}) \to \mathbb{G}(S)$. This isomorphism is induced by the map

$$
\begin{aligned}
&c \mapsto c',\\
&h_i \mapsto y_i,\ i = 1, \dots, q,\\
&p_1^{k_l} \mapsto p_{1,1}\\
&p_1^{(p_l^{j_1-1})} \mapsto p_{1,j_1},\ j_1 = 2, \dots, k_1\\
&p_i^{k_1} \mapsto p_{i,1},\ i = 2, \dots, q,\\
&p_i^{(p_1^{j_i-1})} \mapsto p_{i,j_i},\ j_i = 2, \dots, k_i - 1,\ i = 2, \dots, l\\
&\left(p_i^{k_l-k_i+1}\right)^{p_1^{k_i-1}} \mapsto p_{i,k_i},\ i = 2, \dots, l\\
&x_{i,m_i}^{p_1^{r_i}} \mapsto x_{i,r_i+1,m_i},\ r_i = 0, \dots, k_i - 1,\ i = 1, \dots, l,\ m_i = 1, \dots, v_i\\
&x_{1,m_1}^{p_l^{s}} \mapsto x_{1,s+1,m_1},\ s = 0, \dots, k_1 - 1,\ m_1 = 1, \dots, v_1.
\end{aligned}
$$

In other words, this means that $\pi_1(\mathbb{Y})$ is the reduced centralizer splitting of $\mathbb{G}(S)$. We conclude that $H \simeq \mathbb{G}(S)$. □

Proposition 2. *Let $T = T((v_1, k_1), \dots, (v_l, k_l); p)$ and $S = T((v_1, k_1), \dots, (v_l, k_l); q)$ be two trees of diameter 4. Then there exists r such that $\mathbb{G}(R)$ is a finite index subgroup of both $\mathbb{G}(S)$ and $\mathbb{G}(T)$, where $R = T((v_1, k_1), \dots, (v_l, k_l); r)$. In particular, $\mathbb{G}(S)$ and $\mathbb{G}(T)$ are commensurable.*

Proof. Let $K = k_1 + \dots + k_l$. Since the group $\mathbb{G}(S)$ (the group $\mathbb{G}(T)$ correspondingly) retracts onto the free subgroup F_{K+q} (F_{K+p} correspondingly) generated by pivots and hair vertices, the full preimage of a subgroup of F_{K+q} (of F_{K+p} correspondingly) of finite index I is an index I subgroup of $\mathbb{G}(S)$ ($\mathbb{G}(T)$, correspondingly).

We define subgroups of F_{K+q} and F_{K+p} via covers of bouquet of $K+q$ and $K+p$ circles correspondingly. The subgroup A_q of F_{K+q} is defined as follows. Take $p+K-1$ points $a_1, \dots, a_{p+K-1}$ and for every pivot generator of F_{K+q} add a cycle $(a_1, \dots, a_{p+K-1})$ of length $p+K-1$ labelled by this generator. We complete the obtained graph to a cover of F_{K+q} by adding q loops at every vertex a_i, $i = 1, \dots, p+K-1$. The subgroup A_p of F_{K+p} is defined in a similar fashion.

We note that A_q and A_p are free subgroups of index $p+K-1$ and $q+K-1$ in F_{K+q} and F_{K+p} correspondingly, and both have rank $(q+K-1)(p+K-1)+1$.

Define the subgroups $B_q < \mathbb{G}(S)$ and $B_p < \mathbb{G}(T)$ as the full preimages of the subgroups A_q and A_p correspondingly. We claim that the groups B_q and B_p are isomorphic to $\mathbb{G}(R)$, where $R = T((v_1, k_1), \dots, (v_l, k_l); (q+K-1)(p+K-1)+1-K)$. The proof of this is analogous to the proof of Proposition 1, but easier. It follows from the fact that both B_q and B_p split as fundamental groups of the star of groups with $K = k_1 + \dots + k_l$ leaves. Then the claim holds with $r = (q+K-1)(p+K-1)+1-K$. □

Theorem 5. *Let $T = T((m_1, k_1), \dots, (m_l, k_l); p)$ and $S = T((n_1, r_1), \dots, (n_l, r_l); q)$ be two trees of diameter 4. Suppose that $\frac{m_i}{n_i} = \frac{m_j}{n_j}$ for all $i, j = 1, \dots, l$. Then the group $\mathbb{G}(T)$ is commensurable to $\mathbb{G}(S)$.*

Proof. By Propositions 1 and 2 it suffices to prove the statement in the case when $l \geq 2$, $k_i = r_j = 1$ for $i, j = 1, \dots, l$ and $p = q = 0$, and in the case when $l = 1$, $k_1 = r_1 = 2$ and $p = q = 0$.

Consider the first case, the proof in the second case is analogous. Let $\frac{m_i}{n_i} = \frac{m}{n}$. Consider the homomorphism $f_n : \mathbb{G}(T) \to \mathbb{Z}_n$ induced by the map $c_T \mapsto 1$ and $x \mapsto 0$, where c_T is the center of the tree T and x is any other canonical generator of $\mathbb{G}(T)$. Let $\mathbb{G}_n$ be the kernel of f_n. Similarly, let $f_m : \mathbb{G}(S) \to \mathbb{Z}_m$ be the homomorphism induced by the map $c_S \mapsto 1$ and $y \mapsto 0$, where c_S is the center of the tree S and y is any other canonical generator of $\mathbb{G}(S)$, and let $\mathbb{G}_m$ be the kernel of f_m.

By Bass-Serre theory, it is not difficult to see that $\mathbb{G}_n \simeq \mathbb{G}(T((nm_1, 1), \dots, (nm_l, 1); 0))$ and $\mathbb{G}_m \simeq \mathbb{G}(T((mn_1, 1), \dots, (mn_l, 1); 0))$. Since $nm_i = mn_i$ for all $i = 1, \dots, l$, it follows that $\mathbb{G}_n \simeq \mathbb{G}_m$ and hence $\mathbb{G}(T)$ and $\mathbb{G}(S)$ are commensurable. □

We now turn our attention to the description of minimal elements in the commensurability classes. We prove that the RAAGs defined by paths of length 3 and 4 are commensurable.

Proposition 3. *$\mathbb{G}(P_4)$ is commensurable with $\mathbb{G}(P_5)$.*

Proof. Let $\mathbb{G}(P_4) = \langle x, p_1, c, p_2 \mid [x, p_1], [p_1, c], [c, p_2] \rangle$ and let $\varphi : \mathbb{G}(P_4) \to \mathbb{Z}/(2\mathbb{Z})$ be the homomorphism defined by the map

$$x \to 0 \quad p_1 \to 0 \quad c \to 0 \quad p_2 \to 1$$

Set $H = \ker \varphi$. It is clear that H is an index 2 subgroup of $\mathbb{G}(P_4)$.

Let $\varphi' : \mathbb{G}(P_5) \to \mathbb{Z}/(2\mathbb{Z})$ be the homomorphism defined by the map

$$x_1 \to 0 \quad p_1 \to 1 \quad c \to 0 \quad p_2 \to 1 \quad x_2 \to 0$$

and let $H' = \ker \varphi'$, where x_1 and x_2 are the leaves of P_5 at the pivots p_1 and p_2 correspondingly. It is clear that H' is an index 2 subgroup of $\mathbb{G}(P_5)$.

Straightforward application of Reidemeister-Schreier technique shows that $H = \langle x, p_1, c, p_1^{p_2}, x^{p_2}, p_2^2 \rangle$ and $H' = \langle x_1, p_1^2, c, p_2^2, x_2, p_1 p_2 \rangle$ are isomorphic to $\mathbb{G}(\Delta) = \langle a, b, c, d, e, f \mid [a, b] = 1, [b, c] = 1, [c, d] = 1, [d, e] = 1, [c, f] = 1 \rangle$. □

We deduce the following results.

Theorem 6 (Characterisation of commensurability classes)**.** *Let T and T' be two finite trees of diameter 4, $T = T((d_1, k_1), \dots, (d_l, k_l); q)$ and $T' = T((d'_1, k'_1), \dots, (d'_{l'}, k'_{l'}); q')$. Let $G = \mathbb{G}(T)$ and $G' = \mathbb{G}(T')$. Consider the sets $M = M(T)$, $M' = M(T')$. Then G and G' are commensurable if and only if M and M' are commensurable.*

Proof. It is a consequence of Theorem 4 and Theorem 5. □

Corollary. *The groups $G = \mathbb{G}(P_{m_1, m_2})$ and $G' = \mathbb{G}(P_{n_1, n_2})$ are commensurable if and only if $\frac{m_1}{n_1} = \frac{m_2}{n_2}$.*

Theorem 7 (Minimal RAAG in the commensurability class)**.** *Let $T = T((d_1, k_1), \dots, (d_l, k_l); q)$ be a finite tree of diameter 4. Let $\mathcal{C}(T)$ be the commensurability class of $\mathbb{G}(T)$ and let $M = M(T)$ be as above, so $|M| = l$. Then the minimal RAAG that belongs to $\mathcal{C}(T)$ is either the RAAG defined by the tree $T' = T((d'_1, 1), \dots, (d'_l, 1); 0)$, where $M(T')$ is minimal in the commensurability class of M, if $|M| > 1$, or the RAAG defined by the path of diameter 3, that is $\mathbb{G}(P_4)$, if $|M| = 1$.*

Proof. It is a consequence of Proposition 3 and Theorem 6. □

References

[B72] H. Bass, *The degree of polynomial growth of finitely generated nilpotent groups*, Proc. London Math. Soc. 3, 4 (1972), 603-614.

[BJN09] J. Behrstock, T. Januszkiewicz, W Neumann, *Commensurability and QI classification of free products of finitely generated abelian groups*. Proc. Am. Math. Soc. 137, 3 (2009), 811-813.

[BJN10] J. Behrstock, T. Januszkiewicz, W Neumann, *Quasi-isometric classification of some high dimensional right-angled Artin groups*, Groups, Geom. Dynamics 4, 681-692, 2010.

[BN08] J. Behrstock, W Neumann, *Quasi-isometric classification of graph manifold groups*. Duke Math. J. 141, 2 (2008), 217-240.

[BM00] M. Burger, S. Mozes, *Lattices in product of trees*. Inst. Hautes tudes Sci. Publ. Math., 92 (2000), 151-194.

[DW93] P. Deligne and G. D. Mostow, *Commensurabilities among lattices in $PU(1,n)$*, Annals of Mathematics Studies, **132**. Princeton University Press, 1993.

[DK93] G. Duchamp, D. Krob, *Partially Commutative Magnus Transformations*, Internat. J. Algebra Comput. **3** (1993), 15-41.

[D85] M. J. Dunwoody, *The accessibility of finitely presented groups*. Invent. Mathematicae 81, 3 (1985), 449-457.

[Ga16] A. Garrido, *Abstract commensurability and the Gupta-Sidki group*, Groups Geom. Dyn. 10 (2016), 523-543.

[GrW03] R. Grigorchuk, J. Wilson, *A structural property concerning abstract commensurability of subgroups*. J. London Math. Soc. (2), 68(3):671-682, 2003.

[Gr81] M. Gromov, *Groups of polynomial growth and expanding maps*. Inst. Hautes tudes Sci. Publ. Math., 53 (1981), 53-78.

[Gr93] M. Gromov, *Geometric group theory, vol. 2: Asymptotic invariants of infinite groups*. Lond. Math. Soc. Lecture Notes 182, (1993), 1-295.

[H16] J. Huang, *Commensurability of groups quasi-isometric to RAAGs*, arXiv:1603.08586v2

[KPS73] A. Karrass, A. Pietrowski, D. Solitar, *Finite and infinite cyclic extensions of free groups*, J. Australian Math. Soc 16, 04 (1973), 458-466.

[KK13] S. Kim, T. Koberda, *Embedability between right-angled Artin groups*, Geom. Topol. 17 (2013), 493-530.

[Mar73] G. A. Margulis, *Arithmeticity of nonuniform lattices*, Funkcional. Anal. i Prilozen., 7(3):88-89, 1973.

[Ser89] H. Servatius, *Automorphisms of Graph Groups*, J. Algebra, **126** (1989), no. (1), pp. 34-60.

[Si43] C. L. Siegel, *Symplectic geometry*, Amer. J. Math., 65:1-86, 1943.

[St68] J.R. Stallings, *On torsion-free groups with infinitely many ends* Ann. Math. (1968), 312-334.

[Wh10] K. Whyte, *Coarse bundles*, arXiv:1006.3347.

[Wi96] D. Wise, *Non-positively curved squared complexes aperiodic tilings and non-residually finite groups*, Princeton University, 1996.

Ikerbasque - Basque Foundation for Science and Matematika Saila, UPV/EHU, Sarriena s/n, 48940, Leioa - Bizkaia, Spain

E-mail address: montsecasals@gmail.com

Ikerbasque - Basque Foundation for Science and Matematika Saila, UPV/EHU, Sarriena s/n, 48940, Leioa - Bizkaia, Spain

E-mail address: ilya.kazachkov@gmail.com

Matematika Saila, UPV/EHU, Sarriena s/n, 48940, Leioa - Bizkaia, Spain, and Russian foreign trade academy, 4a Pudovkin Street, 119285, Moscow, Russia

E-mail address: zakhar.sasha@gmail.com